



# Definition of Favourable Conservation Status for lowland calcareous grassland

Defining Favourable Conservation Status Project

Richard G. Jefferson

Second edition published March 2024

First published November 2020

[www.gov.uk/natural-england](http://www.gov.uk/natural-england)

NATURAL  
ENGLAND

# Acknowledgements

---

I would like to thank the following people for their contributions to the production of this document:

- Cath Shellswell at Plantlife, Helen Doherty at SNH and Stuart Smith at NRW.
- Dan Abrahams, Andy Brown, Christina Cork, Jonathan Cox, Jon Curson, Kate Fagan, Cat Hudson, Michael Knight, Sally Mousley, Clare Pinches, Dave Stone, and the Defining Favourable Conservation Status team at Natural England.

# Executive summary

---

This document sets out Natural England's view on favourable conservation status for lowland calcareous grassland in England.

Favourable conservation status is the minimum threshold at which we can be confident that the habitat, and its associated species, are thriving in England and are expected to continue to thrive sustainably in the future.

This definition has been produced following the Natural England approach to defining favourable conservation status described in the guidance document [Defining Favourable Conservation Status in England](#).

Section 1 of this document describes the habitat covered by this definition and its ecosystem context.

Section 2 specifies the units used to describe the three favourable conservation status parameters. These are:

- Natural range and distribution (where the habitat occurs).
- Extent (how much habitat there is).
- The structure and function attributes (habitat quality).

Section 3 outlines the evidence considered when developing the definition. This definition is based on the best available evidence on the ecology of lowland calcareous grassland. The evidence covers the current situation, historical changes and possible future changes.

Section 4 sets out the conclusions on the favourable values, which is the value for each of the three parameters when the habitat has achieved favourable conservation status.

This document does not include any action planning, or describe actions, to achieve or maintain favourable conservation status. These will be presented separately, for example within strategy documents.

## Summary definition of favourable conservation status

Lowland calcareous grassland is a widespread but localised habitat of high biodiversity value occurring throughout the English lowlands and along the upland fringe wherever there are suitable calcareous soils. It is normally maintained through grazing by livestock. The extent of this type of grassland has declined significantly over the last 100 years although its overall range has remained stable. Its vulnerability to climate change has been assessed as low and, like other semi-natural grasslands, it provides important ecosystem services.

The habitat will achieve favourable conservation status when the structure and function attributes set out in section 4.3 are met over 95% of the favourable area. This includes

attributes relating to floristic composition, sward structure, soil nutrient and pH status, grazing management and parcel size and connectivity. In particular, to achieve favourable conservation status the vegetation should be broadly typical of the relevant plant communities and their species composition. There should be some bare ground and variation in sward height including scattered scrub of various age classes. The soils should have properties typical of the habitat notably low soil P and a pH in the range 6.5 to 8.5. The habitat should be grazed by livestock usually at a density of 0.15 to 0.6 LU/ha/year, depending on site productivity and conservation objectives. There should be at least some contiguous or connected areas of suitable semi-natural habitat. All species partially or wholly dependent on this habitat should be Least Concern, when assessed using IUCN criteria.

Favourable status will require an increase in the current extent of the habitat by 149,000 ha (approximately 390 % above the baseline of about 38,687 ha) throughout its range and maintenance of its current range and distribution.

# Contents

---

Definition of Favourable Conservation Status for lowland calcareous grassland .....	1
Acknowledgements .....	2
Executive summary .....	3
Contents .....	5
About the Defining Favourable Conservation Status project .....	6
1. Habitat definition and ecosystem context .....	7
2. Units and attributes.....	11
3. Evidence.....	12
4. Conclusions .....	27
References .....	30
Appendix 1: Constant and positive indicator species for lowland calcareous grassland.	37

# About the Defining Favourable Conservation Status project

---

Natural England's Defining Favourable Conservation Status (DFCS) project is defining the minimum threshold at which habitats and species in England can be considered to be thriving. Our Favourable Conservation Status (FCS) definitions are based on ecological evidence and the expertise of specialists.

Through setting our ambition and aspiration for species and habitats, our definitions will inform decision making and actions to achieve and sustain thriving wildlife.

Our FCS definitions will be embedded into delivery of the 25 Year Environment Plan, through the Nature Recovery Network, biodiversity net gain and environmental land management schemes (ELMS).

Conservation bodies will use them to inform their work, including management planning for the land they own. Businesses will have a clear understanding of how their work impacts nature recovery and how they can help contribute to achieving thriving nature.

By considering the evidence for FCS, decisions will be more confident and strategic, with an understanding of their contribution to, or impact on, the national ambition.

# 1. Habitat definition and ecosystem context

---

## 1.1 Habitat definition

The definition of this habitat embraces the enclosed and unenclosed calcareous grasslands found throughout the UK's lowlands and along the upland fringe but normally below approximately 400 m. Many habitat patches are confined to steep valley slopes, escarpments, and coastal cliffs and headlands. More rarely, they occur on relatively level ground, notably in the East Anglian Breckland and on Salisbury Plain. The habitat is usually managed by extensive sheep or cattle grazing or is more rarely maintained by rabbit grazing. Unenclosed swards in the uplands above the moor wall and managed as free-range rough grazing, sometimes in association with unenclosed tracts of other upland habitats, are not covered by this definition and fall within the upland calcareous grassland habitat.

Lowland calcareous grasslands occur throughout England on mostly shallow, free-draining, infertile lime-rich soils over chalk and limestone bedrock. They are occasionally found on other base-rich substrates such as basic igneous rocks and calcareous glacial drift deposits.

The lowland calcareous grassland priority habitat comprises ten National Vegetation Classification types (CG1-CG10) (Rodwell 1992) but only lowland examples of CG9 and CG10 are included within the definition as this is a lowland habitat as defined and described above. The ten types are as follows:

CG1 *Festuca ovina* – *Carlina vulgaris* grassland

CG2 *Festuca ovina* – *Avenula pratensis* grassland

CG3 *Bromus erectus* grassland

CG4 *Brachypodium pinnatum* grassland

CG5 *Bromus erectus* – *Brachypodium pinnatum* grassland

CG6 *Avenula pubescens* grassland

CG7 *Festuca ovina* – *Hieracium pilosella* – *Thymus praecox/pulegioides* grassland

CG8 *Sesleria albicans* – *Scabiosa columbaria* grassland

CG9 *Sesleria albicans* – *Galium sternerii* grassland

CG10 *Festuca ovina*-*Agrostis capillaris*-*Thymus praecox* grassland

This differs from the definition of the Habitats Directive Annex 1 type known as *H6210 Semi-natural dry grasslands and scrubland facies: on calcareous substrates (Festuco-Brometalia)*, in the following ways:

- Both lowland and upland CG9 is included in H6210.
- CG10 is excluded.

It should also be noted that H6211 Semi-natural dry grasslands 'important orchid rich sites' is now included within H6210.

**Other sources:** Jefferson and others 2014; Rodwell and others 2007.

## 1.2 Habitat status

Lowland calcareous grassland is a S41 Priority Habitat in England reflecting its high conservation value. The habitat also supports a large number of priority species - see Section 1.3 Ecosystem context for further details.

The European Red List of Habitats (Janssen and others 2016) classified the equivalent to the lowland calcareous grassland (EUNIS type – Semi-dry perennial calcareous grassland - 1.2a) as Vulnerable (VU) primarily due to declines suffered over the last 50 years. Specifically, this means a greater than or equal to 30% but less than 50% decline over the last 50 years; a likely future decline greater than or equal to 30% but less than 50% and historic losses since about 1750 of greater than or equal to 50% but less than 70%.

## 1.3 Ecosystem context

Vegetation similar to that of present-day calcareous grassland was probably present in England in the early Holocene where it would probably have occurred on steep slopes, thin soils or exposed terrain which prevented tree growth and/or maintained by native herbivores in glades amongst woodland (for example, Poschlod & WallisDeVries 2002). The limited evidence available supports this hypothesis (Bush & Flenley 1987). The clearance of forest in the Neolithic period onwards for pastoral agriculture would have led to an expansion of such grassland on suitable calcareous substrates.

Subsequently, these unproductive grasslands have had a long history of pastoral agricultural management, being typically used for sheep or cattle rearing.

The occurrence of lowland calcareous grassland in the landscape is largely determined by the occurrence of suitable calcareous geology and soils (rendzina or calcareous brown earths with a pH in the range 6.5 to 8.5) with individual species composition especially influenced by climate and microclimatic factors. The species composition is also affected by geographical location, altitude, topography and aspect. That said, there are examples of where the former quarrying of chalk or limestone, the creation of man-made structures using calcareous rock (for example, railway embankments) or even the deposition of

calcareous industrial wastes (Lee & Greenwood 1976) has resulted in the development, by natural colonisation, of calcareous grassland with very close affinities to 'ancient' calcareous grassland (Jefferson & Usher 1986). In some cases, these dry grasslands may have developed in areas beyond their natural range due to the exposure of limestone substrates that may previously have been covered by drift or overburden.

Calcareous grasslands typically form mosaics with scrub and woodland communities (such as NVC types W8 *Fraxinus excelsior-Acer campestre -Mercurialis perennis* woodland, W9 *Fraxinus excelsior-Sorbus aucuparia-Mercurialis perennis* woodland, W12 *Fagus sylvatica- Mercurialis perennis* woodland, W13 *Taxus baccata* woodland and W21 *Crataegus monogyna-Hedera helix* scrub).

These mosaics and transitions are mediated by past and current management, which is usually related to grazing pressure, including cessation. Where soil type changes, due to drift or head deposits over calcareous rocks, then mosaics may be formed with neutral and acid grasslands and dwarf-shrub heath. Spring lines at the base of chalk or limestone slopes or escarpments may give rise to mire or wetland communities such as M22 *Juncus subnodulosus-Cirsium palustre* fen-meadow or M10 *Carex dioica-Pinguicula vulgaris* mire.

Where acid sands patchily overlie calcareous rocks such as in Breckland, then complex mosaics of calcareous and acid grassland may occur, and sometimes both may occur together with mosaics of heather (*Calluna vulgaris*)-dominated heathland. At a smaller scale, calcareous grasslands may support microhabitats such as short or long turf, grass tussocks, patches of bare ground, rock outcrops and scrub of different age classes.

In general terms, mosaics involving other habitats are more highly valued in conservation assessments (for example, Jefferson and others 2014) and provide an important resource for species (especially invertebrates) that may require a range of habitats or habitat patches and often larger tracts of semi-natural vegetation to complete their life cycle (Webb and others 2009). They may also be indicative of areas or landscapes where natural processes relating to soils and hydrology have been less modified by intensive land management (Mainstone and others 2018).

As with other types of semi-natural grassland (Bullock and others 2011), lowland calcareous grassland can provide ecosystem services including nutrient capture, carbon storage, pollination, pest control, cultural benefits to society and genetic resources.

## **GB and European context**

Comparable grasslands are widespread through western and central Europe, with the parched types such as CG1 having affinities to calcareous grasslands in the sub-Mediterranean zone. Variations in calcareous grassland vegetation across Europe are largely related to differences in climate and soils with a clear geographic sequence from more neutral swards on less drought-prone soils in the Atlantic zone through to steppic grasslands and steppes on the very arid soils in the extreme continental climate of the Urals to south-east Europe.

However, one aspect of British calcareous grasslands that is distinctive is the fading of the more strictly Continental contingent of herbaceous species towards the north of England where lowland calcareous grassland of the Bromion phytosociological alliance (most British calcareous grassland types are placed within this alliance) reaches its north-western European limit. Here, in the Derbyshire Dales and the Pennines, calcareous swards conforming to the CG2d *Dicranum* sub-community of CG2 *Festuca ovina-Avenula pratensis* grassland represents a transition to more acidic swards of the Violion alliance with reduced frequency of typical calcicolous species.

Calcareous grasslands, which were once widespread in Europe, are now a scarce and threatened habitat. There are no overall estimates available for the extent of this habitat type in Europe as a whole. In the rest of the UK, calcareous grasslands occur widely on suitable limestone substrates in Wales but are much more restricted in Scotland and Northern Ireland (Rodwell and others 2007).

The Annex 1 habitat H6210 occurs in 25 European countries within 7 biogeographical regions. The UK has the 6<sup>th</sup> largest area of H6210 in Europe and the 2<sup>nd</sup> largest area within the Atlantic Biogeographical region (European Union in prep).

**Other sources:** Barr 1997; Crofts & Jefferson 1999; Duffey and others 1974; Mortimer and others 2000; Preston & Hill 1997; Rodwell 1991,1992; Smith 1980; UK Biodiversity Group 1998; Veen and others 2009.

## 2. Units and attributes

---

### 2.1 Natural range and distribution

There is a close link between this habitat with geology and soils thus National Character Area (NCA) is considered as an appropriate metric. Figure 2 shows the distribution of the habitat by NCA and highlights the close link between distribution of the habitat and NCAs dominated by calcareous geology such as the Cotswolds, South Downs, Yorkshire Wolds & the Morecambe Bay limestones NCAs.

### 2.2 Extent

Hectare.

### 2.3 Structure and function attributes

Grazing is a key factor in maintaining the structure and function of lowland calcareous grassland assuming the abiotic components of the ecosystem are suitable (soils including nutrient status) and there are no adverse external impacts such as point source or diffuse pollution.

The presence of a moderate number of sheep, cattle or hardy ponies promotes the development and maintenance of the habitat and its component biodiversity by reducing the number of competitive, robust species and allowing other small pioneer plants to thrive. Some sites benefit from rabbit grazing combined with other forms of physical disturbance for maintaining the necessary open conditions, bare ground and short swards required for certain taxa especially invertebrates and lower plants.

The detail of the structure and function attributes are listed in Section 4.3

### 3. Evidence

All blocks of evidence are assigned one of three confidence levels (High, Moderate, Low), based on the quality of the evidence, its applicability and the level of agreement.

The matrix in Figure 1 is used to assess the confidence level assigned to blocks of evidence. White = High confidence; Light blue = Moderate confidence and Dark blue = Low confidence.

Limited evidence Strong agreement	Medium evidence Strong agreement	Robust evidence Strong agreement
Limited evidence Medium agreement	Medium evidence Medium agreement	Robust evidence Medium agreement
Limited evidence Weak agreement	Medium evidence Weak agreement	Robust evidence Weak agreement

**Figure 1 Matrix used to assign confidence to blocks of evidence (after IPCC 2010).**

Quality of evidence is defined as follows:

- Robust evidence is that which has been reported in peer-reviewed literature, or other reputable literature, from well-designed experiments, surveys or inventories that shows signs of being applicable generally.
- Medium evidence is that reported from well-designed experiments, surveys or inventories but from only one or a small number of sites, with uncertainty over its more general applicability, or is correlational or circumstantial evidence.
- Limited evidence includes ‘expert opinion’, based on knowledge of ecological factors that plausibly suggest an effect, but there is no circumstantial or direct evidence available.

Agreement is defined as follows:


- Strong agreement is consensus across the literature and amongst those with expertise on the habitat or species.
- Medium agreement is common consensus across the literature and amongst experts but there are some differing papers or reports and/or some differences of opinion.

- Weak agreement is little consensus across the literature and amongst experts and, possibly, many different findings and/or opinions.


## 3.1 Current situation

### Natural range and distribution

The most up to date information on range/distribution is provided by Rodwell and others 2007 (see Figure 3 below). The number of occupied 10 km squares in the UK is 545 of which 399 (73%) are in England. England supports the greatest proportion of lowland calcareous grassland in the UK at around 95%. There remains the possibility that new sites may be discovered which could extend the current known distribution. However, the discovery of any significant new tract of calcareous grassland is unlikely as calcareous grassland is one of the best-known habitat types in England and has been subject to much survey effort (Blackstock and others 1999). However, the range is ultimately dependent on the availability of suitable substrates although the grassland can sometimes occur beyond the range of suitable natural substrates where artificial lime-rich wastes have been colonised by the suite of species typical of the habitat.



**Figure 2 Distribution of lowland calcareous grassland by National Character Area.**



**Figure 3 Distribution of lowland calcareous grassland by 10 km square**

### **Extent**

It has been estimated that there is 38,687 ha of lowland calcareous grassland in England (Bullock and others 2011). An earlier assessment (Robertson & Jefferson 2000) was very similar at 38,450 ha.

Note: This habitat area estimate may differ from the national Priority Habitat area used for the England Biodiversity Indicator Report. The process of mapping inventories in England rounds areas up to parcel level, is based on old survey data so does not necessarily

reflect recent changes and takes a broader definition of 'grassland' – including partially degraded and less species-rich grassland. Overall, this leads to an overestimate of Priority Habitat cover in England. While extent figures may differ, the England national inventory maps (used in England Biodiversity Indicator reporting) are a good indication of the location of known high quality grassland sites.

## **Patch size and connectivity**

A recent analysis (Bullock and others 2011) showed that of the calcareous grassland sites in the Priority Habitat Inventory, 73% were less than 5 ha with only 7% over 20 ha. The Habitat Network Mapping gives a mean patch size value of 10.7 ha, a median of 1.9 ha with 90% of sites falling below 14.5 ha. As these areas of lowland calcareous grassland priority habitat have a skewed size distribution (that is, many are very small), the median (derived by ranking the patches in size order and taking the size of the middle rank) is a better measure of the 'typical' size than the mean.

## **Quality of habitat patches**

There is much less information on patch quality. Common standards monitoring (CSM) data for SSSIs provides an assessment of condition. In 2018, 45% (approximately 17,100 ha) of the lowland calcareous grassland resource was in favourable condition with 55% unfavourable. A sample survey of non-statutory grasslands undertaken in 2004-2005 (Hewins and others 2005) showed that only 28% of lowland calcareous grasslands sites were in favourable condition. A repeat of the 2004-2005 sample survey was undertaken in 2017 & 2018 but as yet the results are not available.

Historically, (that is prior to the 1970s before the advent of large-scale habitat surveys and assessment methodologies) there is little information on patch quality.

## **Threatened species**

A large number of rare and threatened plants, lichens and fungi are associated with lowland calcareous grassland.

### **Vascular plants**

Around two-thirds of threatened and near threatened grassland vascular plants species are associated with calcareous grassland. The total includes some that are S41 Priority Species such as rare spring sedge *Carex ericetorum*, basil thyme *Clinopodium acinos* and musk orchid *Herminium monorchis* or are listed on Annex II of the Habitats Directive for example, early gentian *Gentianella anglica*.

Threatened or near threatened species listed in the England vascular plant Red List (Stroh and others 2014) include purple milk-vetch *Astragalus danicus*, hoary rock-rose *Helianthemum oelandicum* subsp. *incanum* orchids such as burnt orchid *Neottinea ustulata* and man orchid *Orchis anthropophorum* and devil's bit-scabious *Succisa pratensis*. Some

of these species are range-restricted (hoary rock-rose) whilst others remain widespread (devil's-bit scabious) but all have suffered significant declines in recent years.

Many of these species are specialists of open, short swards normally maintained by grazing animals, both wild and domestic. This guild includes, for example, pasque flower *Pulsatilla vulgaris* which has suffered a 34% decline in Area of Occupancy over the Red List assessment period in part due to a decline in short swards as a result of lack of grazing (Walker & Pinches 2011) and is now regarded as “vulnerable to extinction” (VU). Another example is field fleawort *Tephrosia integrifolia* (classed as Vulnerable) which suffered a 45% decline at 22 sites in southern England over the period 1964-1967 to 2014-2016. A key factor resulting in unsuitable habitat conditions was under grazing or periods of management neglect (Stroh and others 2017).

By contrast, there are a number of threatened species that are associated with tall-herb/scrub margin vegetation/scrub especially on the northern limestones. Such species include bloody cranesbill *Geranium sanguineum*, mezereon *Daphne mezereum* and common valerian *Valeriana officinalis*.

All the component calcareous grassland NVC types are plant species-rich (approximately 20-40+ species per 4 m<sup>2</sup>) when under favourable management.

## **Bryophytes**

There are more bryophytes associated with calcareous grassland, in particular chalk grassland, than any other grassland type in Britain (Porley & Hodgetts 2005). Many are strict calcicoles, including *Ctenidium molluscum*, *Campyliadelphus chrysophyllus*, *Homalothecium lutescens* and *Hypnum lacunosum*. On cool and damp north-facing calcareous slopes, bryophytes can be so abundant that their cover exceeds that of the vascular plants. One community of bryophytes on grazed north-facing chalk slopes on a small number of calcareous grassland sites is unusually rich in liverworts, including *Scapania aspera* and *Porella arboris-vitae*, and has been named the southern hepatic mat. This is considered to be of European conservation significance particular as it appears to be confined to Great Britain with most sites on the English chalk (Porley & Rose 2001).

On steep, south-facing calcareous slopes the combination of warmth from the sun, erosion and grazing by livestock may create a short, thin, parched and disturbed turf with many bare patches where thermophilous (warmth-loving) bryophytes thrive, including some threatened species. In undisturbed calcareous grassland, yellow meadow ants *Lasius flavus* may create large mounds, and these anthills provide a specific habitat which is favoured by certain bryophytes, including the uncommon and declining *Rhodobryum roseum*.

Four S41 bryophytes occur in short and open calcareous turf, namely the liverwort *Cephaloziella baumgartneri* and the mosses *Acaulon triquetrum*, *Weissia condensata* and *Weissia sterilis*.

## Lichens

In addition to the lichen-dominated types of calcareous grassland (the *Cladonia* spp. and *Ditricum flexicaule-Diploschistes scuposus* sub-communities of CG7 *Festuca ovian-Hieracium pilosella-Thymus praecox/pulegiodes* grassland – CG7b and CG7c) calcareous grassland provides an extremely important habitat for the overall diversity of lichens including rare and threatened species (Gilbert 1993) of which two S41 species *Fulgensia fulgens* and scaly Breck lichen *Squamarina lentigera* are examples. The lichen flora may be associated with bare ground, ant hills, chalk pebbles or flints within the grassland. Grassland rich in terricolous lichens (that is, growing on soil or on the ground) is largely restricted to the Cretaceous chalk, especially in southern England. A key habitat requirement for this suite of lichens is the maintenance of open conditions either through disturbance generated by grazing, especially rabbits, or some other form of disturbance which can be anthropogenic (military training for example). Natural processes such as landslips or extreme environmental conditions (for example, a combination of very shallow infertile soil coupled with high temperatures on south-facing slopes).

## Fungi

Calcareous grassland along with other types of semi-natural acid and neutral grassland is considered to have international importance for its fungal assemblages especially of waxcap fungi and certain other fungi groups such as the earth tongues and Clavarioid fungi (Bosanquet and others 2018).

## Vertebrates

A range of vertebrates utilise lowland calcareous grassland and associated habitats, especially scrub, for breeding, feeding or roosting. These include species of conservation concern such as stone curlew, grey partridge and adder.

## Invertebrates

The invertebrate fauna is very diverse and there are many specialist species associated with particular plant species or particular microhabitats found on calcareous bedrock and soils such as bare ground, rocky outcrops and more locally, springs and seepages.

The Pantheon database lists invertebrate species according to habitat traits, though these are rarely if ever exclusive to a particular habitat. A search of Pantheon (Webb and others 2017) using 'calcareous substrate' coupled with a search for eight specific calcareous grassland plant species produced a list of 206 invertebrate species illustrating the utility of Pantheon for conservation assessments and extraction of invertebrate data for specific habitats.

Lowland calcareous grassland supports a large number of rare and threatened invertebrate species. A high proportion of these are species with pronouncedly southern distributions in Britain many of which, at British latitudes, appear to require sunny

situations on well-drained soils. These conditions are often particularly well met on lowland calcareous grassland.

Buglife (2018) list 18 S41 priority species as part of a longer list of 382 notable species associated with the habitat. The former include: leaf beetles such as the Hazel pot beetle *Cryptocephalus coryli*; butterflies such as the silver spotted skipper *Hesperia comma* and Adonis blue *Lysandra bellargus*; the wart-biter cricket *Decticus verrucivorus*; the phantom hoverfly *Doros profuges* and several moths including the bordered gothic *Heliophobus reticulata*, pale shining brown *Polia bombycina* and the four spotted *Tyta luctuosa*.

As a final addendum, research over the last 40 years has demonstrated that the yellow meadow ant *Lasius flavus* is a keystone species in lowland calcareous grassland. It builds long-lasting mounds which considerably increase habitat heterogeneity and overall plant and animal species richness. The flora and fauna of the mounds is distinct from that of the surrounding grassland and often provides a refuge for annual and biennial plants that may have fewer niches in the adjacent grassland. For example, the uncommon winter annual wall whitlow-grass *Draba muralis* is often found on ant hills in calcareous grassland.

Ant hills are also an important resource for other faunal species such as green woodpecker *Picus viridis*. The density, sizes and shapes of the mounds can be used to interpret grassland history. A more detailed but accessible summary is provided by King (2006).

Fagan and others (2010) also found that the ant *Myrmica sabuleti* abundance was significantly positively correlated with age of calcareous grassland restoration sites. The species was a good indicator for calcareous grassland restoration success.

**Confidence:** Moderate - High

### 3.2 Historical variation in the above parameters

All types of lowland semi-natural grassland, including calcareous grasslands, have suffered large losses in the 20th century in lowland England & Wales with an estimated 97% loss over the period 1930 to 1984 (Fuller 1987). More recently a 47% loss has been reported between 1960 and 2013 on sites known to have supported semi-natural grassland but SSSIs were found to have retained more grassland (91%), compared with non-protected sites (27%), thus highlighting their effectiveness as a means of protecting semi-natural grasslands (Ridding, Redhead & Pywell 2015).

Over the last 90 years, lowland calcareous grasslands have been degraded or lost due to agricultural intensification. These changes have involved conversion of grassland to arable, by ploughing, reseeding and treatment with fertilisers and herbicides.

In addition to direct loss, significant areas have ceased to be used for grazing and/or have experienced a loss of their rabbit populations, resulting, in many cases, in sites losing plant diversity, with fewer open areas for specialist species and/or larger areas becoming

dominated by scrub. Abandonment of management can be as damaging for this type of grassland as agricultural intensification. Increasing fragmentation and isolation of sites has been and is an ongoing threat, with many lowland calcareous grassland sites now small in extent (see section 3.1).

## **Loss & fragmentation**

Island biogeography theory predicts that the decreasing extent and fragmentation of semi-natural grasslands over the last 50-100 years will ultimately result in losses of species from remaining areas of grassland habitat. Long transient times in response to decreasing habitat area and increasing isolation due to fragmentation may cause the present plant (and animal) species distribution to reflect the historical rather than the present landscape configuration. Hence, current species populations are possibly not yet in equilibrium with the current landscape configuration, but rather reflect that of the past.

Such 'extinction debt' has been demonstrated for semi-natural grasslands in continental Europe (Krauss and others 2010; Saar and others 2012) but to date, not for England or the UK. However, the principles outlined in Lawton and others (2010) of 'better, bigger, more and joined' should be applied if they are to be conserved effectively. Also, practically, at an individual site level, species populations on small or isolated patches are undoubtedly at a greater risk of extinction for a number of reasons: increased ratio of edge to area increases their susceptibility to external factors such as fertiliser drift; increased probability that stochastic events such as drought and fire will cause extinction across the entire site; tendency to be at greater risk of deterioration in habitat quality over time and their dependence on migrants from larger habitat patches to maintain viable populations.

There is ecological evidence of the negative effects of fragmentation and isolation on the populations of some of the characteristic vascular plants of this and other semi-natural grassland habitats through, for example, genetic erosion (for example, see Matthies and others 2004).

## **Natural range and distribution**

The natural range of calcareous grassland is ultimately dependent on the availability of suitable calcareous substrates. It has probably not radically changed over the last 100 years. Some new areas of suitable substrate outside of the natural range have arisen due to limestone quarrying activities and waste generated by specific industries. These areas can support calcareous grassland due to natural colonisation by the suite of species typical of the habitat. However, within the range the distribution, extent and size of individual sites has markedly changed as detailed below.

## **Extent**

Significant areas of lowland calcareous grassland have been lost during the last 60 years, though it is rarely possible to provide accurate figures. In addition, losses of calcareous grassland to arable cultivation also predate the post-war agricultural revolution and Keymer & Leach (1990) and Jefferson and others (2014) provide some figures of losses

for selected geographical areas from the early 1800s to the 1990s. For example, there was an estimated loss of 76% of the area of chalk grassland in Dorset between 1793 and 1811. At a smaller scale, but typical of more widespread losses, there was an 88% reduction in the area of chalk grassland on Chanctonbury Hill, West Sussex between 1947 and 1991 (Robertson & Jefferson 2002).

An analysis by Burnside and others 2003 found that there was a 42% loss of calcareous grassland in the western South Downs between 1971 and 1991 largely due to agricultural intensification, especially conversion of grassland to arable. Over the same time the number and density of patches of grassland also declined - a 72% decline in patch number.

### **Patch size and connectivity**

An assessment of patch size based on data from the late 1960s for chalk grassland (a subset of calcareous grassland) (Keymer & Leach 1990) showed that of 1,225 documented sites or fragments 76% were between 2 and 20 ha in size with only around 12% exceeding 40 ha. A later analysis, although not fully comparable (Bullock and others 2011) showed that of the calcareous grassland sites in the Priority Habitat Inventory, 73% were less than 5 ha with only 7% over 20 ha. This indicates that in addition to an overall reduction in extent, there was also a decline in size of remaining patches.

### **Quality of habitat patches**

Detailed data on patch quality (such as site condition assessments) is only available for a limited period (last 15-20 years) and mostly for SSSIs (see section 3.1). Patch quality is likely to have declined over the last 50-100 years although no detailed studies are available (see 3.3 for more detail).

The known ecological impacts are summarised below, but it is important to emphasise that actual detailed evidence to support these impacts is rarely available:-

- Reduction in the extent and variety of plant and animal-rich assemblages that occur on calcareous substrates managed by grazing
- Reduction in the species-richness of the calcareous grassland plant communities through impacts such as atmospheric nitrogen deposition and fertiliser overspread and at least the potential for species extinctions through 'extinction debt'
- Reduction in the populations of specialist species associated with the habitat due to impacts such as grazing abandonment, insufficient habitat area to maintain viable populations and eutrophication of soils – see some examples given under ecosystem context (section 1.3)

**Confidence:** Low - Moderate

**Other sources:** Barr 1997; Bobbink and others 1998; Jefferson & Pinches 2018; Tilman and others 1994; Van den Berg and others 2011.

## 3.3 The future for the habitat and its conservation

### Current pressures and threats

The main pressure and threat is a lack of, reduced or inappropriate grazing. Other significant pressures include negative impacts from nutrient enrichment (nitrogen) from atmospheric sources (Van den Berg and others 2011), agricultural intensification including ploughing and grassland improvement via the use of fertilisers. Vulnerability to climate change has been assessed as low but recent work (Basto and others 2018) has highlighted potential negative impacts on vegetation composition and soil seed banks from long-term drought which is predicted to occur with climate change. Basto and others 2018 thus highlight a potential increased climate change vulnerability over the longer term. Therefore, the vulnerability assessment should be regularly reviewed as the results of further research are appraised.

### Natural range and distribution

There is no existing evidence to suggest that there will be any change in range and distribution in the future and maintenance of the current range and distribution should conserve the floristic and faunal diversity associated with the habitat. However, given the evidence from European studies showing extinction debt for grasslands (Section 3.2) then this conclusion may need to be revised. It is expected to be relatively robust to the direct threats posed by climate change, although the climate space of some of its component species is projected to change. The potential range of the habitat is probably unlikely to change greatly over the next 50-100 years but there may be changes in the distribution of certain component species, including range expansions. In addition, with regards to southern calcareous grasslands, potential colonisation by species occurring in northern France in particular, will be constrained by the Channel therefore there is a risk of vacant niches.

### Extent

There is little information on which to make an assessment of the habitat area and site size required for the future maintenance of biodiversity. Some expansion would be very desirable to make up for historic losses and ensure sustainability of threatened species populations such as those mentioned in section 3.1 above.

There are 3 possible approaches to deriving a figure for the habitat area required for the future maintenance of biological diversity or Favourable Conservation Status.

1) Use the indicative area of habitat potential from Natural England's Habitat Network Mapping project. This indicates that potentially 1,500,000 ha have appropriate soils for the development of lowland calcareous grassland.

2) Use the guidance within Defining Favourable Conservation Status in England (Mousley, Van Vliet & Cork 2023). This method uses a "rule-of-thumb" to derive a figure for restoring

a proportion of the historical loss of the habitat. When applied to the restoration of lowland calcareous grassland, this indicates an ambition to restore 90-100% of the historical loss (based on the current status of the habitat as Vulnerable, a high number of associated threatened species and the potential for restoration being 'good'). Assuming a loss of 97% of the habitat (and therefore the current extent is 3% of the historical extent) this would require a minimum increase in area of approximately 1,126,000 ha.

3) Use data produced by the NE National Habitat Network Mapping project. This would indicate an increase of approximately 149,000 ha (390% increase). This is based on the figure required to create a connected network of habitat incorporating existing habitat patches.

It is recommended that the area specified in 3) is adopted. Of the remaining two area expansion targets, 1) represents the full potential for the habitat and favourable conservation status is not considered to require the full potential. Option 2) is very ambitious and would represent over a 3000% increase on the current area. It is considered that option 2) is too ambitious and this level of increase is unlikely to be required to achieve favourable conservation status.

Option 3) is relatively ambitious. However, it is considered that the extent is justifiable given our knowledge of historical losses of calcareous grassland (Keymer & Leach 1990) and the likely negative impacts of decreased patch size and connectivity (see section 3.2). In addition, in terms of suitable habitats for restoration, there are likely to be sufficient areas of land that have the potential for restoration. Also, Barr (1997) calculated that the area of land in England that could potentially support the habitat in terms of suitable substrate amounted to 26.5 M hectares of which the current extent of calcareous grassland is approximately 1.5%. Barr (1997) also calculated that 1,140,000 ha may have at one time been calcareous grassland and was in a land use that could potentially revert. The current extent is around 3% of this estimate.

## **Patch size and connectivity**

Little is known on what constitutes a viable patch size for this habitat but seemingly, provided management is appropriate, patches as small as 0.5 ha may be viable in the long-term. This is the minimum size for qualification as an SSSI (Jefferson and others 2014).

A focus on conserving larger patch sizes and ensuring greater connectivity is clearly important (for example, Lawton and others 2010). However, as pointed out by Reisch and others (2017), a comprehensive conservation approach should also consider the genetic diversity of calcareous grassland plant species. They concluded that populations in smaller grassland fragments may, depending on historical landscape configuration, substantially contribute to the genetic variation of the plant species even under conditions of habitat deterioration. These populations in small fragments should therefore be included in strategies to conserve calcareous grassland.

Certain animals require larger patch sizes, although precise details are often lacking. One species where some information is available is the Marsh Fritillary a species associated with both purple moor-grass and rush pastures and lowland calcareous grassland. Marsh fritillary populations function on a landscape-scale. They are often highly cyclical with large fluctuations in population size, making them prone to local extinction, but this characteristic also allows the butterfly to colonise new sites in good years as well as patches of less suitable habitat. The butterfly persists in areas where large networks of suitable habitat exist, with groups of local populations being connected by occasional dispersal, known as metapopulations. To achieve long-term population stability, the butterfly requires an extensive network of well-connected habitat patches where the food plant, devil's-bit scabious (*Succisa pratensis*), is abundant. At a site level, optimal management usually involves extensive grazing regimes using cattle at around 0.4 livestock units per hectare. Research has shown that an area of between 80 ha and 142 ha per 1,600 ha (that is, 5-9% of a landscape) is required to achieve persistence for 100 years, depending on the spatial location of the habitat (Bulman and others 2007).

### Quality of habitat patches

Although the evidence base is limited, there is a likelihood that patch quality and species diversity of calcareous grassland sites has probably declined over the last 90 years, at least in some geographical areas, due to a range of impacts including nutrient enrichment and management changes (see 3.2). The lack of available data on these changes makes it challenging to decide on what constitutes an appropriate baseline. However, there is definitely scope for analysis of historic calcareous grassland datasets to examine this issue in more detail.

A particular example of a shifting baseline, although not calcareous grassland, are data from the long-running Rothamsted Park Grass neutral grassland plots showed a shifting baseline of species richness in plots with the baseline dropping by 15 species m<sup>2</sup> over a 130-year period (Dodd and others 1994).

Should the CSM favourable condition baseline plus species richness and species composition variables from the National Vegetation Classification and other unpublished vegetation data (Rodwell 1992) be used as measures of quality and diversity then, assuming the pressures of nutrient enrichment and lack of management are reduced over time, then quality and diversity of patches can be maintained or even enhanced.

**Confidence:** Low.

## 3.4 Constraints to expansion or restoration

The most common cause of unfavourable condition of existing sites is a lack of or inappropriate grazing. This can be readily addressed although practical reasons relating to the availability of grazing animals or the need for associated infrastructure (fencing, water supply, access etc) can hinder progress. In some cases, nutrient enrichment from local point sources can result in unfavourable conditions which can take several decades to

reverse depending on the level of intervention. Calcareous grasslands are less prone than some other types of semi-natural grassland to the impact of atmospheric nitrogen deposition as they are P-limited systems.

Restoration and creation of new calcareous grasslands has been the focus of activity as a result of Biodiversity Action planning and agri-environment schemes over the last 20 years. The targeting and methodology of grassland habitat restoration is now well understood due to a considerable body of research effort (see Pywell and others 2013).

There are many examples of successful restoration for example, Wilson and others 2013) although the timescales for such grasslands to resemble 'ancient' examples, at least in terms of their floristic composition, may take up to at least a century (Gibson & Brown 1991; Redhead and others 2014; Wagner and others 2019).

The best examples of restoration though are where very infertile, exposed calcareous substrates border or are surrounded by existing calcareous grassland. These appear to readily re-colonise naturally over time (Wilson and others 2013).

From an invertebrate perspective, Woodcock and others (2015) found that when the aim is to create new grasslands on ex-arable, establishing plant communities typical of species-rich grassland is a prerequisite for ensuring the success of grassland creation. This can be achieved by the use of appropriate seed-mixtures or the use of green hay / brush-harvested seed (natural seeding). However, landscape factors, primarily the proximity of existing semi-natural grassland, plays an important role in determining whether invertebrates successfully colonise sites irrespective of their floral composition. Immediate proximity of existing sites is likely to be very important for colonisation by less mobile species such as some species of grasshoppers and crickets.

Allison and others (2016) provided evidence that for the conservation of specialist macro-moth species restricted to calcareous grassland, agri-environment scheme (AES) interventions are most effective if positioned close to the habitat.

## **Natural range and distribution**

Natural range/distribution would be largely unaffected unless former quarrying created opportunities for restoration in areas beyond the natural distribution of calcareous geology and soils.

## **Extent**

In theory, there are no immovable constraints to increasing the area of the habitat subject to having a favourable policy and legislative environment and the necessary financial resources, the latter most likely via agri-environment payments.

The recommendations in Lawton and others 2010 are highly relevant here in that this report advocates i) the improvement of the quality of existing sites, ii) increasing the size of

existing sites and iii) enhancing the connectivity of existing sites by way of habitat restoration and creation.

### **Patch size and connectivity**

Patch size and connectivity can readily be increased through habitat creation and restoration.

### **Quality of habitat patches**

Given adequate resourcing, there should be no hindrance to achieving structural and functional requirements on sites though for newly created sites, as noted above, it may take many decades to achieve the quality attributes. In areas where calcareous grassland occurs in areas dominated by arable agriculture, there would be benefits in a general shift towards more mixed farming systems which might possibly be facilitated by provision of targeted incentive schemes. This would provide more opportunities for sustainable grazing management.

The impact of lowland calcareous grassland restoration and expansion will be beneficial for grassland species assemblages, priority species and for certain species requiring larger areas of habitat such as stone curlew.

**Confidence:** High.

# 4. Conclusions

---

## 4.1 Favourable range and distribution

The current range (see section 3.1) is likely to be favourable for future maintenance of the habitat. The favourable area will fall within the current range (including historic and current occurrences) but will ultimately include the areas of restored habitat that constitute Favourable Conservation status for the habitat. The range could be monitored using a combination of climate, topographical and soil/substrate parameters.

## 4.2 Favourable extent

The favourable area is set at approximately 188,000 ha which is the current area increased by 149,000 ha. This would involve restoration largely from semi-improved grassland, arable and dense scrub. The rationale for this ambition is set out in section 3.2. Temporal changes in the area could be monitored by a combination of field-based sample-based monitoring and earth observation methods.

## 4.3 Favourable structure and function attributes

### Structure attributes

- Typical vegetation community and species composition at desired frequency and cover (Appendix 1 lists typical and positive indicator plant species for the habitat).
- Natural pattern of vegetation zonation/transitions/mosaics.
- Low cover of undesirable species.
- The presence of some bare ground for regeneration niches and supporting habitat for specialist Invertebrates, vascular plants, bryophytes, lichens and fungi.
- The presence of rock outcrops, chalk/limestone pebbles and flints especially in relation to bryophytes and lichens.
- Micro and macro topographic heterogeneity.
- The presence of ant hills of the yellow meadow ant in sites within the species natural range (see section 3.1).
- Vegetation heterogeneity (including scattered scrub of different age classes) and suitable 'floweriness' (nectar/pollen resources) to benefit fauna especially invertebrates (See Webb and others 2009).

## Function attributes

- Properties of the underlying geology (solid and drift) – calcareous rocks and drift deposits, including equivalent artificial alkaline substrates.
- Properties of the underlying soil types within typical values for the habitat, including: structure, bulk density, total carbon, pH, exchangeable soil calcium (Ca), exchangeable soil acidity, soil nutrient status and fungal: bacterial ratio. For this feature, soil P index should typically be index 0 (less than 9 mg per litre). However, P indices of 1 or 2 have been measured on sites which may have been cultivated in the past.
- Supporting off-site habitat for example, contiguous or connected areas of suitable habitats.\*
- Functional connectivity with the wider landscape.\*
- Concentrations and deposition of air pollutants at or below the site-relevant Critical Load or Level values.
- Suitable grazing management by livestock appropriate to deliver conservation objectives, usually within the range 0.15 to 0.6 LU/ha/year depending on site productivity and conservation objectives.
- Presence and activity of rabbits at sustainable levels (principally creation of bare ground and short turf) to deliver habitat and species objectives at selected sites.
- Biotic and abiotic processes which create and maintain bare ground and early successional communities (for example, slope processes – soil creep, solifluction).

\* These two attributes are included as there is some evidence for the benefits of larger habitat patches and general connectivity of sites both between calcareous grasslands but also other habitat types. The evidence is summarised in Lawton and others 2010 and further elucidated in the Natural England Nature Networks Handbook (Crick and others 2020).

Note that the structure and function attributes apply to the habitat and it may be that certain species of conservation concern (see section 3.1) may require additional attributes to those detailed above to meet FCS.

## Quality of habitat patches

At least 95% of the favourable area of the habitat meets the structure and function requirements as described above.

## Threatened species

All species partially or wholly dependent on this habitat should be Least Concern, when assessed using IUCN criteria (or considered to be Least Concern if not formally assessed), as regards to this habitat.

**Sources:** Bullock and others 2011; Crick and others 2020; Critchley and others 2002; Crofts & Jefferson 1999; Duffey and others 1974; King 2006; Kirkham and others 2003; Lawton and others 2010; Robertson & Jefferson 2000; Rodwell 1992; Webb and others 2009.

# References

---

- Allison, J., Duffield, S. J., Van Noordwijk, C. G. E., Morecroft, M. D., Marrs, R. H., Saccheri, I. J., and Hodgson, J. A. 2016. Spatial targeting of habitat creation has the potential to improve agri-environment scheme outcomes for macro-moths. *Journal of Applied Ecology*, 53, 1814–1822.
- Barr, C. J. (Ed.). 1997. *Current status and prospects for key habitats in England. Part 2 Calcareous grassland landscapes*. Department of the Environment, Transport & the Regions, London.
- Basto, S., Ken Thompson, K., Grime, J. P., Fridley, J. D., Calhim, S., Askew, A. P., and Rees, M. 2018. Severe effects of long-term drought on calcareous grassland seed banks. *Climate & Atmospheric Science*, 1, 1-7. <https://doi.org/10.1038/s41612-017-0007-3> (Accessed 22 March 2024).
- Blackstock, T. H., Rimes, C. A., Stevens, D. P., Jefferson, R. G., Robertson, H. J., Mackintosh, J., and Hopkins, J. J. 1999. The extent of semi-natural grassland communities in lowland England and Wales: a review of conservation surveys 1978-96. *Grass and Forage Science*, 54, 1-18.
- Bobbink, R., Hornung, M., and Roelofs, J. G. M. 1998. The effects of air-borne nitrogen pollutants on species diversity in natural and semi-natural European vegetation. *Journal of Ecology*, 86, 717–738.
- Bosanquet, S., Ainsworth, M., Cooch, S., Genney, D., and Wilkins, T. 2018. *Guidelines for the Selection of Biological SSSIs. Part 2: Detailed Guidelines for Habitats and Species Groups. Chapter 14: Non-lichenised fungi*. Joint Nature Conservation Committee, Peterborough.
- Buglife. 2018. Lowland calcareous Grassland. <https://www.buglife.org.uk/advice-and-publications/advice-on-managing-bap-habitats/lowland-calcareous-grassland> (Accessed 22 March 2024).
- Bullock, J. M., Jefferson, R. G., Blackstock, T. H., Pakeman, R. J., Emmett, B. A., Pywell, R. J., Grime, J. P., and Silvertown, J. W. 2011. Chapter 6: Semi-natural grasslands. *In: The UK National Ecosystem Assessment Technical Report*. UK National Ecosystem Assessment, UNEP-WCMC, Cambridge.
- Bulman, C. R., Wilson, R. J., Holt, A. R., Bravo, L. G., Early, R. I., Warren, M. S., and Thomas, C. D. 2007. Minimum viable metapopulation size, extinction debt and the conservation of a declining species. *Ecological Applications*, 17: 1460-1473.
- Burnside, N. G., Smith, R. F., and Waite, S. 2003. Recent historical land use change on the South Downs, United Kingdom. *Environmental Conservation*, 30, 52-60.

Bush, M. B., and Flenley, J. R. 1987. The age of British chalk grassland. *Nature*, 329, 434-436.

Crick, H. Q. P., Crosher, I. E., Mainstone, C. P., Taylor S. D., Wharton, A., Langford, P., Larwood, J., Lusardi, J., Appleton, D., Brotherton, P. N. M., Duffield, S. J., and MacGregor N. A. 2020. *Nature Networks Evidence Handbook. Natural England Research Report NERR081*. Natural England, York.

Critchley, C. N. R., Chambers, B. J., Fowbert, J. A., Sanderson, R. A., Bhogal, A., and Rose, S. C. 2002. Association between lowland grassland plant communities and soil properties. *Biological Conservation*, 105, 199-215.

Crofts, A., and Jefferson, R. G. 1999. *The Lowland Grassland Management Handbook*. English Nature & The Wildlife Trusts, Peterborough.

Dodd, M. E., Silvertown, J., McConway, K., Potts, J., and Crawley, M. 1994. Stability in the plant communities of the Park Grass Experiment: the relationships between species richness, soil pH and biomass variability. *Philosophical Transactions of the Royal Society Series B*, 346, 185-193.

Dolman, P. M., Panter, C. J., and Mossman, H. L. 2010. *Securing Biodiversity in Breckland: Guidance for Conservation and Research. First Report of the Breckland Biodiversity Audit*. University of East Anglia, Norwich.

Duffey, E., Morris, M. G., Sheail, J., Ward, L. K., Wells, D. A., and Wells, T. C. E. 1974. *Grassland ecology and wildlife management*. London, Chapman and Hall.

European Union in prep. EU Habitat Action Plan. *Action plan to maintain and restore the habitat type 6210 Semi-natural dry grasslands and scrubland facies on calcareous substrates (Festuco-Brometalia) (\*important orchid sites) to favourable conservation status*.

Fagan, K. C., Pywell, R. F, Bullock, J. M., and Marrs, R. H. 2010. Are Ants Useful Indicators of Restoration Success in Temperate Grasslands? *Restoration Ecology*, 18, 373-379.

Fuller, R. M. 1987. The changing extent and conservation interest of lowland grasslands in England and Wales: a review of grassland surveys 1930-1984. *Biological Conservation*, 40, 281-300.

Gibson, C. W. D., and Brown, V. K. 1991. The nature and rate of development of calcareous grassland in Southern Britain. *Biological Conservation*, 58, 297-316.

Gilbert, O. L. 1993. The Lichens of chalk grassland. *The Lichenologist*, 25,379-414.

Hewins, E. 2012. *A survey of selected agri-environment grassland and heathland creation and restoration sites: Part 1. Natural England Commissioned Report, NECR107*. Natural England, Sheffield.

Hewins, E. J., Pinches, C., Arnold, J., Lush, M., Robertson, H., and Escott, S. 2005. *The condition of lowland BAP priority grasslands: results from a sample survey of non-statutory stands in England. English Nature Research Reports, No 636.* English Nature, Peterborough.

Jackson, D. L., and MCleod, C. R. 2002. *Handbook on the UK status of EC Habitats Directive interest features: provisional data on the UK distribution and extent of Annex I habitats and the UK distribution and population size of Annex II species. JNCC Report 312.* Joint Nature Conservation Committee, Peterborough.

Janssen, J. A. M., Rodwell, J.S., García Criado, M., Gubbay, T., Haynes, A., Nieto, N., Sanders, F., Landucci, J., Loidi, A., Ssymank, T., Tahvanainen, M., Valderrabano, A., Acosta, M., Aronsson, G., Arts, F., Attorre, E., Bergmeier, R.-J., Bijlsma, F., Bioret, C., Biță-Nicolae, I... Valachovič, M. 2016. *European Red List of Habitats. Part 2. Terrestrial and freshwater habitats.* Publications Office, European Union.

Jefferson, R. G., and Pinches, C. 2018. Article 17 submission to the Joint Nature Conservation Committee. Unpublished database.

Jefferson, R. G., Smith, S. L. N., and Mackintosh, E. J. 2014. *Guidelines for the Selection of Biological SSSIs. Part 2: Detailed Guidelines for Habitats and Species Groups. Chapter 3 Lowland Grasslands.* Joint Nature Conservation Committee, Peterborough.

Jefferson, R. G., and Usher, M. B. 1986. Ecological succession and the evaluation of non-climax communities. In: M.B. USHER, ed. *Wildlife Conservation Evaluation*, pp 70-91. London: Chapman & Hall.

Keymer, R. J., and Leach, S. J. 1990. Calcareous grassland – a limited resource in Britain. In: Hillier, S. H., Walton, D. W. H, and Wells, D. A., (eds.). *Calcareous Grasslands – Ecology and Management. Proceeding of a British Ecological Society and Nature Conservancy Council Symposium, 1987*, pp 11-17. Bluntisham Books, Huntingdon.

King, T. J. 2006. The value of ant hills in grasslands. *British Wildlife*, 17, 392-397.

Kirkham, F. W., Mole, A., Gardner, S. M., and Wilson, D. W. 2003. *Review of Stocking Levels Recommended for Semi-natural Lowland Grasslands. Countryside Council for Wales (CCW) Contract Science Report No. 596.* CCW, Bangor.

Krauss, J., Bommarco, R., Guardiola, M., Heikkinen, R. K., Helm, A., Kuussaari, M., Lindborg, R., Öckinger, E., Pärtel, M., Pino, J., Pöyry, J., Raatikainen, K. M., Sang, A., Stefanescu, C., Teder, T., Zobel, M., and Steffan-Dewenter, I. 2010. Habitat fragmentation causes immediate and time-delayed biodiversity loss at different trophic levels. *Ecology Letters*, 13 : 597–605.

Lawton, J. H., Brotherton, P. N. M., Brown, V. K., Elphick, C., Fitter, A. H., Forshaw, J., Haddow, R. W., Hilborne, S., Leafe, R. N., Mace, G. M., Southgate, M. P., Sutherland, W. J., Tew, T. E., Varley, J., and Wynne, G. R. 2010. *Making Space for Nature: a Review of*

*England's Wildlife Sites and Ecological Network. Report submitted to the Secretary of State, the Department for Environment, Food and Rural Affairs on 16 September 2010.* Defra, London.

Lee, A., and Greenwood, B. 1976. The colonisation by plants of calcareous wastes from the salt and alkali industry in Cheshire, England. *Biological Conservation*, 10: 131-149.

Mainstone, C., Jefferson, R. G., Diack, I., Alonso, I., Crowle, A., Rees, S., Goldberg, E., Webb, J., Drewitt, A., Taylor, I., Cox, J., Edgar, P., and Walsh, K. 2018. *Generating more integrated biodiversity objectives – rationale, principles and practice. Natural England Research Report No. 071 + lowland grassland Appendix C.*

Matthies, D., Brauer, I., Maiborn, W., and Tschardtke, T. 2004. Population size and the risk of local extinction: empirical evidence from rare plants. *Oikos*, 105: 481-488.

Mortimer, S. R., Turner, A. J., Brown, V. K., Fuller, R. J., Good, J. E. G., Bell, S. A., Stevens, P. A., Norris, D., Bayfield, N., and Ward, L. K. 2000. *The nature conservation value of scrub in Britain. JNCC Report 308.* Joint Nature Conservation Committee, Peterborough.

Mousley, S., Van Vliet, W., and Cork, C. 2023. *Defining favourable conservation status in England: Natural England approach.* Natural England Evidence Information Note EIN062. York: Natural England.

Natural England. 2008. *State of the Natural Environment 2008.* Natural England, Sheffield. <http://publications.naturalengland.org.uk/publication/31043>. (Accessed 22 March 2024).

Natural England, and RSPB. 2014. *Climate Change Adaptation Manual: Evidence to support nature conservation in a changing climate. Natural England Commissioned Research Report no. 546.*

Pinches, C. E., and Jefferson, R. G. 2011. Grassland data at the UK scale: development and applications. *In: Blackstock, T. H., Howe, E., Rothwell, J. P., Duigan, C. A., and Jones, P. S. Proceedings of a memorial conference for Dr David Paul Stevens, 1958-2007. Grassland Ecologist and conservationist. CCW Staff Science Report No 10/03/05.* Countryside Council for Wales, Bangor.

Porley, R. D., and Hodgetts, N. 2005. *Mosses and Liverworts.* Collins New Naturalist Series, No. 97 Collins, London.

Porley, R. D., and Rose, F. 2001. The characterization and status of the southern hepatic mat, *Scapanietum asperae* Rose & Porley, on the English Chalk. *Journal of Bryology*, 23, 195-204.

Poschlod, P., and Wallis Devries, M. F. 2002. The historical and socioeconomic perspective of calcareous grasslands—lessons from the distant and recent past. *Biological Conservation*, 104:361–376.

- Preston, C. D., and Hill, M. O. 1997. The geographical relationships of British and Irish vascular plants. *Botanical Journal of the Linnean Society*, 124: 1-120.
- Pywell, R. F., Woodcock, B., Tallowin, J. R. B., Mortimer, S. R., and Bullock, J. M. 2012. Restoring species-rich grassland: principles and techniques. *In: Peel, S., Chesterton, C., Cooke, A., Jefferson, R. G., Martin, D., Smith, B., Smith, S., and Tallowin, J. R. B., (Eds.). Restoring diverse grasslands: What can be achieved where, and what will it do for us? Aspects of Applied Biology*, 115, 11-21.
- Ratcliffe, D. A. 1977. *A Nature Conservation Review*. Cambridge University Press, Cambridge.
- Redhead, J. W., Sheail, J., Bullock, J. M., Ferreruella, A., Walker, K. J., and Pywell, R. F. 2014. The natural regeneration of calcareous grassland at a landscape scale: 150 years of plant community re-assembly on Salisbury Plain, UK. *Applied Vegetation Science*, 17, 408–418.
- Ridding, L. E., Redhead, J. W., and Pywell, R. F. 2015. Fate of semi-natural grassland in England between 1960 and 2013: a test of national conservation policy. *Global Ecology and Conservation*, 4, 516-525.
- Robertson, H. J., and Jefferson, R. G. 2000. *Monitoring the condition of lowland grassland SSSIs. Volume I: English Nature's rapid assessment system. English Nature Research Reports 315, Volume I*. English Nature, Peterborough.
- Robertson, H. J., and Jefferson, R. G. 2002. Nature and Farming in Britain. *In: D.L. Jackson, and L.L. Jackson, (Eds.)., The Farm as Natural Habitat: Reconnecting food systems with ecosystems*. Island Press, Washington D.C.
- Reisch, C., Schmidkonz, S., Meier, K., Schöppllein, Q., Meyer, C., Hums, C., Putz, C., and Schmid, C. 2017. Genetic diversity of calcareous grassland plant species depends on historical landscape configuration. *BMC Ecol*, 17, 19. <https://doi.org/10.1186/s12898-017-0129-9>. (Accessed 22 March 2024).
- Robertson, H. J., Jefferson, R. G., Wilson, P. J., Reed, M. E., McDonnell, E., and Wheeler, B. 2002. *Monitoring grassland Biodiversity Action Plan targets: condition and restoration assessment methodologies for non-statutory grasslands. Interim Report*. English Nature, Peterborough.
- Rodwell, J. S. (Ed.). 1991. *British Plant Communities Volume 2: Mires and heaths*. Cambridge: Cambridge University Press.
- Rodwell, J. S. (Ed.). 1992. *British Plant Communities Volume 3: Grasslands and Montane Communities*. Cambridge University Press, Cambridge.
- Rodwell, J. S., Morgan, V., Jefferson, R. G., and Moss, D. 2007. *The European context of British lowland grasslands. JNCC Report 394*. Joint Nature Conservation Committee, Peterborough.

Saar, L., Takkis, K., Pärtel, M., and Helm, A. 2012. Which plant traits predict species loss in calcareous grasslands with extinction debt? *Diversity and Distributions*, 18: 808-817.

Smith, C. J. 1980. *Ecology of the English Chalk*. Academic Press, London.

Stevens, C. J., Smart, S. M., Henrys, P. A., Maskell, L. C., Walker, K. J., Preston, C. D., Crowe, A., Rowe, E. C., Gowing, D. J., and Emmett, B. A. 2011. *Collation of evidence of nitrogen impacts on vegetation in relation to UK biodiversity objectives*. JNCC Report, No. 447.

Stroh, P. A., Leach, S. J., August, T. A., Walker, K. J., Pearman, D. A., Rumsey, F. J., Harrower, C. A., Fay, M. F., Martin, J. P., Pankhurst, T., Preston, C. D., and Taylor, I. 2014. *A Vascular Plant Red List for England*. Botanical Society for Britain and Ireland, UK.

Stroh, P. A., Pescott, O. L., and Mountford, J. O. 2017. Long-term changes in lowland calcareous grassland plots using *Tephrosia integrifolia* subsp. *integrifolia* as an indicator species. *Plant Ecology*, 218:1269-1281.

Tilman, D., May, R. M., C. L. Lehman, C. L., and Nowak, M.A. 1994. Habitat destruction and the extinction debt. *Nature*, 371: 65–66.

UK Biodiversity Group. 1998. *Tranche 2 Action Plans. Volume II – terrestrial and freshwater habitats*. English Nature, Peterborough, UK.

Van Den Berg, L. J. L., Vergeer, P, Rich, T. C. G., Smart, S. M., Guest, D., and Ashmore, M. R. 2011. Direct and indirect effects of nitrogen deposition on species composition change in calcareous grasslands. *Global Change Biology*, 17: 1871–1883.

Veen, P., Jefferson, R. G., De Smidt, J., and Van Der Straaten, J. (Eds.). 2009. *Grasslands in Europe of High Nature Value*. KNNV Publishing, Zeist, The Netherlands.

Wagner, M., Fagan, K. C., Jefferson, R. G., Marrs, R. H., Mortimer, S. R., Bullock, J. M., and Pywell, R. F. 2019. Species indicators for naturally regenerating and old calcareous grassland in southern England. *Ecological indicators*, 101, 804-812.

Walker, K. J., and Pinches C. E. 2011. Reduced grazing and the decline of *Pulsatilla vulgaris* Mill. (Ranunculaceae) in England, UK. *Biological Conservation*, 144:3098–3105.

Walker, K. J., Stroh, P. A., and Ellis, R. W. 2018. *Threatened Plants of Britain and Ireland*. Botanical Society of Britain and Ireland, Bristol.

Webb, J. R., Drewitt, A. L., and Measures, G. H. 2009. *Managing for species: integrating the needs of England's priority species into habitat management*. Research Report NERR024. Natural England, Sheffield.

Webb, J., Heaver, D., Lott, D., Dean, H. J., Van Breda, J., Curson, J., Harvey, M., Gurney, M., Roy, D. B., Van Breda, A., Drake, M., Alexander, K. N. A., and Foster, G. 2017.

Pantheon - database version 3.7.6. (online) Available at: <http://www.brc.ac.uk/pantheon/> (Accessed: 26 September 2017).

Wilson, P., Wheeler, B., Reed, M., and Strange, A. 2013. *A survey of selected agri-environment grassland and heathland creation and restoration sites.: Part 2. Natural England Commissioned Report, NECR107*. Natural England, Sheffield.

Woodcock, B., Ball, S., Amy, S., Edwards, M., Redhead, J., Mountford, J. O., Gregory, S., Duffield, S., MacGregor, N., and Pywell, R. F. 2015. *Grassland Restoration for Ecological Networks: Evaluating the relative importance of site and landscape characteristics for invertebrate communities in grasslands restored through agri-environment schemes. Final report for the GREEN project (LM0450/ECM6418)*. Defra, London.  
<http://randd.defra.gov.uk/Default.aspx?Menu=Menu&Module=More&Location=None&ProjectID=19417&FromSearch=Y&Publisher=1&SearchText=LM0450&SortString=ProjectCode&SortOrder=Asc&Paging=10#Description>. (Accessed 22 March 2024).

# Appendix 1: Constant and positive indicator species for lowland calcareous grassland

---

## Constant species from component National Vegetation Classification types

*Avenula pratensis*, *Avenula pubescens*, ***Brachypodium rupestre*** (formerly ***pinnatum***) , *Briza media*, ***Bromopsis erecta***, ***Campanula rotundifolia***, *Carex flacca*, ***Carlina vulgaris***, ***Cirsium acaule***, *Ctenidium molluscum*, *Dactylis glomerata*, *Festuca ovina*, *Festuca rubra*, ***Galium sternerii***, ***Helianthemum nummularium***, *Hypnum cupressiforme*, *Koeleria macrantha*, ***Leontodon hispidus***, ***Linum catharticum***, ***Lotus corniculatus***, ***Pilosella officinarum***, *Plantago lanceolata*, ***Poterium sanguisorba***, *Pseudoscleropodium purum*, ***Scabiosa columbaria***, ***Sesleria caerulea***, *Taraxacum* agg, ***Thymus polytrichus/pulegiodes***, *Viola riviniana*.

## Typical species indicating favourable condition (from Robertson & Jefferson 2000 and additions from Robertson and others 2002)

Those in bold above plus:

*Agrimonia eupatoria*, *Anthyllis vulnaria*, *Arenaria serpyllifolia*, *Asperula cyanachica*, *Astragalus danicus*, *Betonica officinalis*, *Blackstonia perfoliata*, *Campanula glomerata*, *Centaurea nigra* agg, *Centaurea scabiosa*, *Centaureum erythraea*, *Cladonia* spp, *Clinopodium acinos*, *Clinopodium vulgare*, *Erigeron acris*, *Euphrasia* spp, *Filipendula vulgaris*, *Fragaria vesca*, *Gallium verum*, *Gentianella* spp, *Geranium sanguineum*, *Helianthemum appeninum*, *Helianthemum oelandicum*, *Hippocrepis comosa*, *Hypericum pulchrum*, *Knautia arvensis*, *Lathyrus pratensis*, *Leontodon saxatilis*, *Leucanthemum vulgare*, *Myosotis ramossisimia*, *Onobrychis viciifolia*, *Ononis repens/spinose*, *Orchidaceae* spp, *Origanum vulgare*, *Phyteuma tenerum*, *Pimpinella* spp, *Plantago media*, *Polygala* spp, *Primula veris*, *Scilla* spp, *Sedum* spp, *Serratula tinctoria*, *Succisa pratensis*, *Teucrium scorodonia*, *Tragopogon pratensis*, *Trinia glauca*, *Viola hirta*.

# About Natural England

Natural England is here to secure a healthy natural environment for people to enjoy, where wildlife is protected and England's traditional landscapes are safeguarded for future generations.

## Further Information

This report can be downloaded from the [Natural England Access to Evidence Catalogue](#). For information on Natural England publications or if you require an alternative format, please contact the Natural England Enquiry Service on 0300 060 3900 or email [enquiries@naturalengland.org.uk](mailto:enquiries@naturalengland.org.uk).

## Citation

Jefferson, R.G. 2024. Definition of Favourable Conservation Status for lowland calcareous grassland. Second edition. RP2944. Natural England.

## Copyright

This publication is published by Natural England under the [Open Government Licence v3.0](#) for public sector information. You are encouraged to use, and reuse, information subject to certain conditions.

Natural England photographs are only available for non-commercial purposes. If any other photographs or information such as maps or data cannot be used commercially this will be made clear within the report.

For information regarding the use of maps or data see our guidance on [How to access Natural England's maps and data](#).

Cover image: Chalk grassland, Holies Down SSSI, Berkshire. Des Sussex, Natural England, 2015.

© Natural England 2024

Catalogue code: RP2944

First edition published 26th November 2020.

This edition published March 2024.

