

## AGRICULTURAL LAND CLASSIFICATION

### BLOCK FEN, MEPAL (Proposed Mineral Extraction)

#### 1.0 BACKGROUND

- 1.1 This 131.9 hectare site was inspected between March 1989 and February 1990 in connection with proposals to extract sand and gravel.
- 1.2 A total of 136 auger borings were made over the site on a 100 metre grid basis superimposed on the national grid. These inspections were made using a hand held 120 cm Dutch soil auger and were supplemented by observations from five soil profile pits.
- 1.3 At the time of survey the site was mainly in arable cultivation, typical crops including cereals, sugar beet, potatoes, onions and carrots. A small area (approximately one hectare) towards the southwest corner was under paddock and scrub woodland.

#### 2.0 PHYSICAL FACTORS AFFECTING LAND QUALITY

##### Climate

- 2.1 Area specific climate data has been obtained by interpolating information contained in the 5km grid dataset produced by the Meteorological office, (Met Office, 1989).
- 2.2 The annual average rainfall is approximately 540mm (21.6 inches) which is low by national standards. Soils are likely to be at field capacity for a relatively short period of approximately 88 days between mid March and mid December. During this time the workability of the land is not likely to be greatly impaired due to the relatively free-draining nature of the gravel substrate.
- 2.3 The accumulated temperature for this area is approximately 1457 degrees celsius. This parameter indicates the cumulative build up of warmth available for crop growth, and has an influence on the development of

soil moisture deficits (SMD)\* and susceptibility to drought; the soil moisture deficits for potatoes and wheat are 114mm and 118mm respectively. These figures are slightly higher than average for lowland England.

- 2.4 The site is neither particularly exposed, or frost prone.
- 2.5 There is no overall climatic limitation to the agricultural use of this land, although some of the soils are susceptible to drought.

#### Relief and Drainage

- 2.6 The majority of the site lies between 1 and 2m AOD. However, towards the extreme east of the site the land slopes gently to a minimum altitude of just below 0m AOD adjacent to the ditch. In general terms soils are relatively free draining (predominantly wetness class I). Groundwater levels are controlled by a series of deep ditches and dykes which feed water into a pumped drainage system which discharges into the Hundred Foot Drain. Although localised wetness may occur on the lowlying land adjacent to the ditch at the eastern end of the site, this is unlikely to constitute a long term limitation to land use.

#### Geology and Soils

- 2.7 The published 1:50,000 scale geology map for this area indicates that the site is underlain by Jurassic Ampthill Clay (Geol.Surv, 1980). Over the majority of the site this Clay is overlain by spreads of first

\* SMD represents the balance between rainfall and potential evapotranspiration occurring during the growing season. For ALC purposes the soil moisture deficits developing under a winter wheat and maincrop potato cover are considered. These 'reference' crops have been selected because they are widely grown, and in terms of their susceptibility to drought, are representative of a wide range of crops.

and second terrace river gravels. On the narrow strip of lower lying land to the east however, the Clay is overlain by Nordelph Peat. Field observations support this general description and also indicate that a fairly high degree of variability exists within the terrace soils identified on site, both laterally over short distances and vertically down the soil profile.

- 2.8 In very general terms the terrace soils, comprise sandy loam, sandy silt loam or medium clay loam topsoils over thinly bedded coarse and fine loamy subsoils. Typically subsoils include sandy loam, sandy clay loam, clay loam and loamy sand textures, although these may be interbedded with sand and/or heavier clay-enriched horizons of clay and sandy clay textures at variable depth. The underlying gravel deposit is usually encountered between depths of 80-100cm, although deeper and shallower soil profiles were noted intermittently.
- 2.9 The Nordelph peat, which once formed a mantle over the entire site has now largely wasted from the area of terrace gravel soils. Successive cultivations have progressively incorporated the peat remnant into the underlying mineral soil, resulting in topsoils having enhanced levels of organic matter. Laboratory analyses on a range of typical topsoil textures indicate organic matter levels are in the range 5.2% to 16.4%.\* Enhanced levels of organic matter effectively contribute to the water storage capacity of the soil and help offset any susceptibility to drought stress.
- 2.10 On the low lying land at the eastern end of the site the peat cover remains more intact. Profiles in this area comprise peaty loam topsoils over subsoils of loamy peat which is typically interbedded with clay, silty clay and sandy loam textures. A narrow band of intergrade soils occurs at the junction of the two main soil types.

\* Topsoils which contain between 6%-25% organic matter (on a sliding scale determined by clay content) are defined as "organic".

- 2.11 In general terms the majority of soil profiles are slightly or very slightly stony\* in upper horizons, but tend to become increasingly stony below 60/70cm as the underlying gravel deposit is approached (small areas of moderately to very stony soils were also recorded in random locations). The topsoil stone volumes recorded on site are not sufficient to constitute a limitation to agricultural use, although they will influence the water storage capacity of the soil, and therefore affect its susceptibility to drought.
- 2.12 pH readings were taken throughout the soil profile at intervals across the site. These indicated typical pH values of between 6.5 and 7.0 on the terrace soils, which is within an acceptable range for most crops.
- 2.13 Within the small area of peat derived soils towards the east of the site, field pH readings were consistently below 5.0 from depths of 40cm and below. These acid horizons are formed by the oxidation of pyrite (Ferrous disulphide) which is a stable constituent of some anaerobic marine sediments (MAFF, 1983). When drained, air is allowed to penetrate the soil mass and the pyrite oxidises to form sulphuric acid, which impairs the development of plant roots through the soil profile. Below depths of approximately 40cm this acidity is not easily rectified by normal liming operations and therefore constitutes a limitation to land quality, since plant roots are unable to extract water stored in the soil beneath the acid horizon.
- 2.14 Soil profile pit observations indicate that the terrace soils are porous and free draining (predominantly wetness class I) and are not subject to any significant wetness or workability constraints. Field observations support this assessment and indicate that moderately heavy falls of rain do not significantly delay cultivation or drilling operations, within this area. Towards the extreme east of the site, the small area of lowlying land adjacent the ditch is believed to be affected by slight fluctuations in groundwater levels. However these are not thought to be sufficiently severe to constitute an overriding limitation to land quality.

\* Comprising small and very small subangular flints and rounded pebbles.

2.15 Generally all the soils are non calcareous, although random profiles do exhibit a thin strongly calcareous marl horizon at variable depth. The underlying gravel deposit is mainly strongly calcareous.

### 3.0 AGRICULTURAL LAND CLASSIFICATION

3.1 Land on this site is graded 1, 2 and 3a. A breakdown of ALC grades in hectares and % terms is provided below.

| ALC              | Hectares | Percentage |
|------------------|----------|------------|
| 1                | 21.6     | 16.4       |
| 2                | 47.4     | 35.9       |
| 3a               | 62.0     | 47.0       |
| Non Agricultural | 0.9      | 0.7        |
|                  | <hr/>    | <hr/>      |
| Total            | 131.9    | 100.0      |
|                  | <hr/>    | <hr/>      |

Note: The ALC grading in this report and on the accompanying maps relates to the long term potential of the land without irrigation.

3.2 The majority of the site is limited by droughtiness. The relative severity of this limitation was assessed using the Revised guidelines and criteria for grading agricultural land (MAFF 1989). Crop adjusted available water capacity (AP)\* values were calculated for each sample profile, using maincrop potatoes and winter wheat as reference crops, characteristic of a broad range of arable and horticultural crops.

\* AP is a measure of the quantity of water held in the soil profile which can be taken up by a specified crop. The water storage capacity of soil is influenced by texture, structure, organic matter content and stone content. Where rooting is impeded for chemical or physical reasons, this is also taken into account.

These AP values were then offset against the crop adjusted soil moisture deficit values described in para 2.3 to obtain moisture balance figures for wheat and potatoes. These moisture balance figures indicate the relative degree of the droughtiness limitation and relate directly to ALC grade. A description of the type of land occurring in each grade is provided below:

#### Grade 1

- 3.3 This is mapped in the central and eastern parts of the site. Profiles are typically slightly or very slightly stony in upper horizons and comprise organic sandy loam or medium clay loam topsoils over sandy loam, sandy clay loam, clay loam or clay subsoils which in turn overlie the gravel deposit below 80-90cm. The water storage capacity of land in this grade is high. This derives in part from the enhanced organic matter content of the topsoils and also from the relatively stone free, well-bodied and water retentive subsoils. Drought risk is consequently minimised and the land is capable of producing consistently high yields of a wide range of crops.

#### Grade 2

This occurs in two main situations:

- 3.4 In the area of terrace gravel soils it is mapped where profiles are typically slightly or very slightly stony in upper horizons, and comprise sandy loam or sandy silt loam topsoils over subsoils of sandy loam, sandy clay loam or clay loam, which frequently contain heavier clayey or lighter textured sand or loamy sand horizons at variable depth. Topsoils are typically non organic, although towards the north of the site stonier profiles with organic topsoils are also included in the grade. The water storage capacity of land in this grade is moderately high, and the land is limited by only minor droughtiness constraints.
- 3.5 Grade 2 land is also mapped towards the east of the site where profiles typically comprise peaty loam topsoils overlying subsoils of interbedded peaty loam, sandy loam and silty clay textures. Rooting within the subsoils is likely to be restricted by the presence of

strongly acid horizons, at and below 40cm depth which are not easily removed by normal management operations. The land is consequently limited by droughtiness imperfections which are only partly offset by the high organic matter content of the topsoil.

Grade 3a

- 3.6 This is mapped extensively on site in areas of soils with reduced capacity for soil water storage. This is due to a combination of factors including low levels of topsoil organic matter, higher profile stone volumes, a reduced depth of soil over gravel and a greater proportion of less water retentive loamy sand and sand textures in the upper profile. The land is consequently limited by moderate droughtiness imperfections.
- 3.7 Although individual profiles of better and/or poorer grades were noted within each mapping unit, they occurred too randomly or inextensively to permit separate delineation at the scale shown.

March 1990

Katherine A Jewson  
Resource Planning Group

## STATEMENT OF SITE/SOIL PHYSICAL CHARACTERISTICS

### BLOCK FEN, MEPAL

#### 1.0 SITE PHYSICAL FACTORS

1.1 Details of climate, relief and drainage and geology are included in the full ALC report supplied for this site.

#### 2.0 SOIL PHYSICAL FACTORS

2.1 Field survey observations indicate that a fairly high degree of variability exists in the terrace soils identified on site both laterally over short distances and vertically down the soil profile. For this reason only one soil type has been identified on site and this is described in detail below.

2.2 Towards the extreme east of the site, a narrow ribbon of lowlying land flanking the brook and graded 2 is comprised of peaty textures which are acid at and below 40cm. Some consideration should be given to handling this small area of contrasting soils separately.

#### 3.0 SOIL TYPE 1

Topsoil Texture : sandy loam, sandy silt loam or medium clay loam

Colour : variable, typically 10YR 3/2 & 10YR 3/1

Stone : total stone is typically less than 5% topsoil volume, comprising predominantly of flints and pebbles in the size range 2mm-2cm

Stonier patches (up to 25% topsoil volume) were recorded locally with a similar size distribution.



Depth : although a-typical topsoil depths of between 20 and 50cm were recorded in isolated locations, most topsoils were in the range 30-40cm, with 35cm emerging as the modal depth.

Structure/Porosity: cultivation zone - not applicable.

Boundary : smooth, clear or abrupt lower boundary.

Roots : common fine and very fine roots.

Subsoil Texture : comprise finely bedded coarse and fine loamy soils. Typical textures include sandy loam, sandy clay loam, clay loam, although these may be interbedded with lenses of sand/loamy sand or clay/sandy clay horizons at variable depth.

Colour : In the range 7.5YR 4/6 to 2.5Y 3/6

Stone : In the range 0-70% (for ALC purposes 70% plus stones is assessed as gravel). Size distribution as topsoil.

Depth : variable over short distances. In the range 45-120cm, typically 70-80cm.

Structure : weakly developed coarse subangular blocky or massive. Friable to firm consistence.

Porosity : generally more than .5% biopores (>.5mm).

Boundary : clear, smooth to irregular lower boundary.

Roots : common to 70/80 cm; fewer 70/80cm plus, except in area of acid peaty soils, where rooting is likely to be restricted at shallow depth.

Gravel Deposit : Typically comprises of grit and fine gravel stones which appear in varying quantities in a matrix ranging from sand to clay. Less frequently deposits of stoneless sand are encountered.

Additional Information:

Wetness class Predominantly wetness class I, smaller areas of wetness class II occur locally in more slowly permeable (clayey) subsoil variants.

Calcium carbonate Typically non calcareous in upper horizons may become more calcareous as the gravel deposit is approached. Individual profiles may contain a thin highly calcareous marly horizon at depth.

March 1990

Katherine A Jewson  
Resource Planning Group  
Cambridge RO