

A1
NEWBURY LOCAL PLAN
SITE 7: ALDERMASTON WHARF
AGRICULTURAL LAND CLASSIFICATION
ALC MAP & REPORT
FEBRUARY 1994

**NEWBURY LOCAL PLAN
SITE 7: ALDERMASTON WHARF
AGRICULTURAL LAND CLASSIFICATION REPORT**

1.0 Summary

1.1 ADAS was commissioned by MAFF's Land Use Planning Unit to provide information on land quality on a number of sites in the Newbury District of Berkshire. The work forms part of MAFF's statutory input to the preparation of the Newbury Local Plan.

1.2 Approximately 5 hectares of land relating to site 7, Aldermaston Wharf was surveyed in February 1994. The survey was undertaken at a detailed level of approximately one boring per hectare. A total of 5 soil auger borings and 1 soil pit were assessed in accordance with MAFF's revised guidelines and criteria for grading the quality of agricultural land (MAFF, 1988). These guidelines provide a framework for classifying land according to the extent to which its physical or chemical characteristics impose longterm limitations on its use for agriculture.

1.3 The work was conducted by members of the Resource Planning Team in the Guildford Statutory Group of ADAS.

1.4 At the time of the survey the agricultural landuse on the site was permanent grassland being partly grazed by horses. In addition to this a small area of non-agricultural land has also been mapped which surrounds a farm building and urban development.

1.5 The distribution of grades and subgrades is shown on the attached ALC map and the areas are given in the table below. The map has been drawn at a scale of 1:5,000. It is accurate at this scale, but any enlargement would be misleading. This map supersedes any previous survey information for this site.

Table 1 : Distribution of Grades and Subgrades

<u>Grade</u>	<u>Area (ha)</u>	<u>% of Site</u>	<u>% of Agricultural Area</u>
3b	4.1	80.4	100
Non-Agricultural	0.8	15.7	100%
Urban	<u>0.2</u>	<u>3.9</u>	
Total area of site	5.1	100%	

1.6 Appendix 1 gives a general description of the grades, subgrades and land use categories identified in the survey. The main classes are described in terms of the type of limitation that can occur, the typical cropping range and the expected level and consistency of yield.

1.7 All of the agricultural land surveyed has been classified as moderate quality subgrade 3b due to a significant soil droughtiness limitation. The profiles comprise slightly stony medium clay loam topsoils which become heavier and markedly stonier with depth. The lower subsoils, comprising coarse sand with a high gravel content, appear at moderately shallow depths. The stone content present within these profiles and their relatively shallow depth over gravelly horizons causes the amount of water available for crops to be reduced and therefore gives rise to a significant drought risk.

2.0 Climate

2.1 The climatic criteria are considered first when classifying land as climate can be overriding in the sense that severe limitations will restrict land to low grades irrespective of favourable site or soil conditions.

2.2 The main parameters used in the assessment of an overall climatic limitation are average annual rainfall, as a measure of overall wetness, and accumulated temperature (degree days Jan-June), as a measure of the relative warmth of a locality.

2.3 A detailed assessment of the prevailing climate was made by interpolation from a 5km gridpoint dataset (Met. Office 1989). The details are given in the table below and these show that there is no overall climatic limitation affecting the site. However climatic factors do interact with soil properties to influence soil wetness and droughtiness limitations.

2.4 No local climatic factors such as exposure or frost risk affect the site.

Table 2 : Climatic Interpolation

Grid Reference :	SU603669
Altitude (m) :	55
Accumulated Temperature (days) :	1466
Average Annual Rainfall (mm) :	689
Field Capacity (days) :	147
Moisture Deficit, Wheat (mm) :	111
Moisture Deficit, Potatoes (mm) :	105
Overall Climatic Grade :	1

3.0 Relief

3.1 The site lies at an altitude of 55m. AOD and is relatively flat.

4.0 Geology and Soil

4.1 British Geological Survey (1946), sheet 268, Reading shows the entire site to be underlain by Valley Gravel.

4.2 The soil type for this site, shown on the Soil Survey Map of England and Wales, sheet 268 (SSEW, 1967, 1:63360), comprises the Thames Series in the south and the Sonning 2 Series in the north. The Thames soils are described as 'Stoneless, mainly calcareous clayey soils affected by groundwater. Flat land. Risk of flooding.' while the Sonning are 'Well drained flinty coarse loamy and gravelly soils. Associated with slowly permeable seasonally waterlogged fine loamy over clayey soils with slowly permeable subsoils and slight seasonal waterlogging.' (SSEW, 1983).

Detailed examination of the soils on the site revealed a closer similarity to the Sonning Series than the Thames.

5.0 Agricultural Land Classification

5.1 Table 1 provides the details of the area measurements for each grade and the distribution of each grade is shown on the attached ALC map.

5.2 The location of the soil observation points are shown on the attached sample point map.

5.3 Subgrade 3b

The majority of the soil profiles examined were calcareous and comprised slightly stony (7% flints < 2cm) medium clay loam topsoils over very stony (40% flints < 2cm) heavy clay loam upper subsoils. From a depth of 45-60cm, and occasionally shallower (35cm), the subsoil becomes coarse sand containing 65% flints. There are no signs of soil wetness in any of the horizons so the profiles have been assigned to Wetness Class I. The moderately high stone content in the upper subsoils and the occurrence of coarse sand and gravel lower subsoils cause a significant droughtiness problem. The amount of water available in the profile for plants to extract is markedly reduced and so the crops are likely to suffer drought stress for part or all of the growing season. The land, therefore is restricted to subgrade 3b on soil droughtiness.

5.5 The areas marked as Urban include a building, tarmac drive and parking area.

5.6 The non-agricultural elements of the site comprise a rough area littered with bricks and concrete, a small group of saplings and a private vegetable patch.

ADAS REFERENCE : 0202/024/93
MAFF REFERENCE : EL 02/0297

Resource Planning Team
Guildford Statutory Group
ADAS Reading

APPENDIX I

DESCRIPTION OF THE GRADES AND SUB-GRADES

Grade 1 : Excellent Quality Agricultural Land

Land with no or very minor limitations to agricultural use. A very wide range of agricultural and horticultural crops can be grown and commonly includes top fruit, soft, fruit, salad crops and winter harvested vegetables. Yields are high and less variable than on land of lower quality.

Grade 2 : Very Good Quality Agricultural Land

Land with minor limitations which affect crop yield, cultivations or harvesting. A wide range of agricultural or horticultural crops can usually be grown but on some land on the grade there may be reduced flexibility due to difficulties with the production of the more demanding crops such as winter harvested vegetables and arable root crops. The level of yield is generally high but may be lower or more variable than Grade 1.

Grade 3 : Good To Moderate Quality Agricultural Land

Land with moderate limitations which affect the choice of crops, timing and type of cultivation, harvesting or the level of yield. When more demanding crops are grown yields are generally lower or more variable than on land in grades 1 and 2.

Sub-grade 3A : Good Quality Agricultural Land

Land capable of consistently producing moderate to high yields of a narrow range of arable crops, especially cereals, or moderate yields of a wide range of crops including cereals, grass, oilseed rape, potatoes, sugar beet and the less demanding horticultural crops.

Sub-grade 3B : Moderate Quality Agricultural Land

Land capable of producing moderate yields of a narrow range of crops, principally cereals and grass, or lower yields of a wider range of crops or high yields of grass which can be grazed or harvested over most of the year.

Grade 4 : Poor Quality Agricultural Land

Land with severe limitations which significantly restrict the range of crops and/or the level of yields. It is mainly suited to grass with occasional arable crops (eg. cereals and forage crops) the yields of which are variable. In moist climates, yields of grass may be moderate to high but there may be difficulties in utilisation. the grade also includes very droughty arable land.

Grade 5 : Very Poor Quality Agricultural Land

Land with very severe limitations which restrict use to permanent pasture or rough grazing, except for occasional pioneer forage crops.

Urban

Built-up or 'hard' uses with relatively little potential for a return to agriculture : housing, industry, commerce, education, transport, religious buildings, cemeteries. Also, hard-surfaced sports facilities, permanent caravan sites and vacant land; all types of derelict land, including mineral workings which are only likely to be reclaimed using derelict land grants.

Non-agricultural

'Soft' uses where most of the land could be returned relatively easily to agriculture, including : private parkland, public open spaces, sports fields, allotments and soft-surfaced areas on airports/airfields. Also active mineral workings and refuse tips where restoration conditions to 'soft' after-uses may apply.

Woodland

Includes commercial and non-commercial woodland.

Agricultural Buildings

Includes the normal range of agricultural buildings as well as other relatively permanent structures such as glasshouses. Temporary structures (eg. polythene tunnels erected for lambing) may be ignored.

Open Water

Includes lakes, ponds and rivers as map scale permits.

Land Not Surveyed

Agricultural land which has not been surveyed.

Where the land use includes more than one of the above, eg. buildings in large grounds, and where map scale permits, the cover types may be shown separately. Otherwise, the most extensive cover type will be shown.

APPENDIX II

REFERENCES

- * British Geological Survey (1946), Sheet No.268, Reading, 1:50,000
- * MAFF (1988), Agricultural Land Classification of England And Wales : revised guidelines and criteria for grading the quality of agricultural land.
- * Meteorological Office (1989), Climatological Data for Agricultural Land Classification.
- * Soil Survey of England and Wales (1967), Sheet No.268, Soils of Reading, 1:63360. And accompanying legend (SSEW, 1983).

APPENDIX III

DEFINITION OF SOIL WETNESS CLASSES

Wetness Class I

The soil profile is not wet within 70cm depth for more than 30 days in most years.

Wetness Class II

The soil profile is wet within 70cm depth for 31-90 days in most years or, if there is no slowly permeable layer within 80cm depth, it is wet within 70cm for more than 90 days, but not wet within 40cm depth for more than 30 days in most years.

Wetness Class III

The soil profile is wet within 70cm depth for 91-180 days in most years or, if there is no slowly permeable layer within 80cm depth, it is wet within 70cm for more than 180 days, but only wet within 40cm depth for 31-90 days in most years.

Wetness Class IV

The soil profile is wet within 70cm depth for more than 180 days but not wet within 40cm depth for more than 210 days in most years or, if there is no slowly permeable layer within 80cm depth, it is wet within 40cm depth for 91-210 days in most years.

Wetness Class V

The soil profile is wet within 40cm depth for 211-335 days in most years.

Wetness Class VI

The soil profile is wet within 40cm depth for more than 335 days in most years.

(The number of days is not necessarily a continuous period. 'In most years' is defined as more than 10 out of 20 years.)

APPENDIX IV

SOIL PIT AND SOIL BORING DESCRIPTIONS

- Contents :
- * Soil Abbreviations : Explanatory Note
 - * Soil Pit Descriptions
 - * Database Printout : Boring Level Information
 - * Database Printout : Horizon Level Information

SOIL PROFILE DESCRIPTIONS : EXPLANATORY NOTE

Soil pit and auger boring information collected during ALC fieldwork is held on a database. This has commonly used notations and abbreviations as set out below.

Boring Header Information

1. GRID REF : national grid square and 8 figure grid reference.

2. USE : Land use at the time of survey. The following abbreviations are used.

ARA : Arable WHT : Wheat BAR : Barley CER : Cereals OAT : Oats MZE : Maize OSR : Oilseed rape
BEN : Field Beans BRA : Brassicae POT : Potatoes SBT : Sugar Beet FCD : Fodder Crops LIN : Linseed
FRT : Soft and Top Fruit HRT : Horticultural Crops PGR : Permanent Pasture LEY : Ley Grass RGR : Rough Grazing
SCR : Scrub CFW : Coniferous Woodland DCW : Deciduous Woodland HTH : Heathland BOG : Bog or Marsh
FLW : Fallow PLO : Ploughed SAS : Set aside OTH : Other

3. GRDNT : Gradient as measured by a hand-held optical clinometer.

4. GLEY/SPL : Depth in cm to gleying or slowly permeable layers.

5. AP (WHEAT/POTS) : Crop-adjusted available water capacity.

6. MB (WHEAT/POTS) : Moisture Balance.

7. DRT : Best grade according to soil droughtiness.

8. If any of the following factors are considered significant, an entry of 'Y' will be entered in the relevant column.

MREL : Microrelief limitation FLOOD : Flood risk EROSN : Soil erosion risk EXP : Exposure limitation FROST : Frost
DIST : Disturbed land CHEM : Chemical limitation

9. LIMIT : The main limitation to land quality. The following abbreviations are used.

OC : Overall Climate AE : Aspect EX : Exposure FR : Frost Risk GR : Gradient MR : Microrelief
FL : Flood Risk TX : Topsoil Texture DP : Soil Depth CH : Chemical WE : Wetness WK : Workability
DR : Drought ER : Soil Erosion Risk WD : Combined Soil Wetness/Droughtiness ST : Topsoil Stoniness

Soil Pits and Auger Borings

1. TEXTURE : soil texture classes are denoted by the following abbreviations.

S : Sand LS : Loamy Sand SL : Sandy Loam SZL : Sandy Silt Loam CL : Clay Loam ZCL : Silty Clay Loam
SCL : Sandy Clay Loam C : Clay SC : Sandy Clay ZC : Silty Clay OL : Organic Loam P : Peat SP : Sandy Peat
LP : Loamy Peat PL : Peaty Loam PS : Peaty Sand MZ : Marine Light Silts

For the sand, loamy sand, sandy loam and sandy silt loam classes, the predominant size of sand fraction will be indicated by the use of prefixes.

F : Fine (more than 66% of the sand less than 0.2mm)

M : Medium (less than 66% fine sand and less than 33% coarse sand)

C : Coarse (more than 33% of the sand larger than 0.6mm)

The clay loam and silty clay loam classes will be sub-divided according to the clay content.

M : Medium (<27% clay) H : Heavy (27-35% clay)

2. MOTTLE COL : Mottle colour

3. MOTTLE ABUN : Mottle abundance, expressed as a percentage of the matrix or surface described.

F : few <2% C : common 2-20% M : many 20-40 VM : very many 40% +

4. MOTTLE CONT : Mottle contrast

F : faint - indistinct mottles, evident only on close inspection D : distinct - mottles are readily seen
P : prominent - mottling is conspicuous and one of the outstanding features of the horizon

5. PED. COL : Ped face colour

6. STONE LITH : One of the following is used.

HR : all hard rocks and stones MSST : soft, medium or coarse grained sandstone
SI : soft weathered igneous or metamorphic SLST : soft oolitic or dolimitic limestone
FSST : soft, fine grained sandstone ZR : soft, argillaceous, or silty rocks CH : chalk
GH : gravel with non-porous (hard) stones GS : gravel with porous (soft) stones

Stone contents (> 2cm, > 6cm and total) are given in percentages (by volume).

7. STRUCT : the degree of development, size and shape of soil peds are described using the following notation:

- degree of development WK : weakly developed MD : moderately developed ST : strongly developed

- ped size F : fine M : medium C : coarse VC : very coarse

- ped shape S : single grain M : massive GR : granular AB : angular blocky SAB : sub-angular blocky PR : prismatic
PL : platy

8. CONSIST : Soil consistence is described using the following notation:

L : loose VF : very friable FR : friable FM : firm VM : very firm EM : extremely firm EH : extremely hard

9. SUBS STR : Subsoil structural condition recorded for the purpose of calculating profile droughtiness.

G : good M : moderate P : poor

10. POR : Soil porosity. If a soil horizon has less than 0.5% biopores > 0.5 mm, a 'Y' will appear in this column.

11. IMP : If the profile is impenetrable a 'Y' will appear in this column at the appropriate horizon.

12. SPL : Slowly permeable layer. If the soil horizon is slowly permeable a 'Y' will appear in this column.

13. CALC : If the soil horizon is calcareous, a 'Y' will appear in this column.

14. Other notations

APW : available water capacity (in mm) adjusted for wheat
APP : available water capacity (in mm) adjusted for potatoes
MBW : moisture balance, wheat
MBP : moisture balance, potatoes

SAMPLE NO.	GRID REF	ASPECT USE	--WETNESS--		-WHEAT-		-POTS-		M.REL		EROSN	FROST	CHEM	ALC	COMMENTS
			GRDNT	GLEYSPL	CLASS	GRADE	AP	MB	AP	MB	DRT	FLOOD	EXP	DIST	
1	SU60306700	PGR	1	1	087	-24	088	-17	38					DR 3B	IMP 52+
1P	SU60276687	PGR	1	1	065	-46	066	-39	38					DR 3B	IMP 60+ ALL 3B
3	SU60276690	PGR	1	1	000	0	000	0						DR 3B	IMP 52+
4	SU60206680	PGR	1	1	071	-40	071	-34	38					DR 3B	IMP 46+
5	SU60306680	PGR	1	1	000	0	000	0						DR 4	PROB3B IMP 35+
6	SU60396687	PGR	1	1	058	-53	058	-47	4					DR 4	PROB 3B

SAMPLE	DEPTH	TEXTURE	COLOUR	----MOTTLES-----			PED	----STONES----			STRUCT/	SUBS						
				COL	ABUN	CONT	COL.	GLEY	>2	>6	LITH	TOT	CONSIST	STR	POR	IMP	SPL	CALC
1	0-35	mc1	10YR42 00					0	0	HR	2							
	35-45	hc1	10YR43 00					0	0	HR	2				G			
	45-52	hc1	10YR43 00					0	0	HR	10				G			
1P	0-26	mc1	10YR42 00					2	0	HR	7							Y
	26-40	hc1	10YR53 00					0	0	HR	40	WKMSAB	FR	G				Y
	40-60	cs	10YR72 00					0	0	HR	65	S	L	M				Y
3	0-25	mc1	10YR42 00					0	0	HR	3							Y
	25-48	hc1	10YR53 00					0	0	HR	3				G			Y
	48-52	c	10YR54 00					0	0	HR	10				M			Y
4	0-30	mc1	10YR42 00					0	0	HR	10							Y
	30-46	mc1	10YR62 00					0	0	HR	15				G			Y
5	0-30	mc1	10YR42 00					0	0	HR	3							Y
	30-35	hc1	10YR53 00					0	0	HR	10				G			Y
6	0-33	mc1	10YR42 00					0	0	HR	2							Y

SOIL PIT DESCRIPTION

Site Name : SITE 7,ALDERMASTON WHARF Pit Number : 1P

Grid Reference: SU60276687 Average Annual Rainfall : 689 mm
 Accumulated Temperature : 1466 degree days
 Field Capacity Level : 147 days
 Land Use : Permanent Grass
 Slope and Aspect : degrees

HORIZON	TEXTURE	COLOUR	STONES >2	TOT.STONE	MOTTLES	STRUCTURE
0- 26	MCL	10YR42 00	2	7		
26- 40	HCL	10YR53 00	0	40		WKMSAB
40- 60	CS	10YR72 00	0	65		S

Wetness Grade : 1 Wetness Class : I
 Gleying : cm
 SPL : No SPL

Drought Grade : 3B APW : 065mm MBW : -46 mm
 APP : 066mm MBP : -39 mm

FINAL ALC GRADE : 3B
 MAIN LIMITATION : Droughtiness