

A1
WEST SUSSEX MINERALS PLAN
SITE 15 : WESTERN EXTENSION,
ROCK COMMON
AGRICULTURAL LAND CLASSIFICATION
ALC MAP & REPORT
NOVEMBER 1993

**WEST SUSSEX MINERALS PLAN
SITE 15 : WESTERN EXTENSION, ROCK COMMON
AGRICULTURAL LAND CLASSIFICATION REPORT**

1.0 Summary

1.1 ADAS was commissioned by MAFF's Land Use Planning Unit to provide information on land quality on a number of sites in West Sussex. The work forms part of MAFF's statutory input to the preparation of the West Sussex Minerals Plan.

1.2 Approximately 4 hectares of land relating to site 15, at Rock Common West was surveyed in November 1993. The survey was undertaken at a detailed level of approximately one boring per hectare. A total of 5 soil auger borings and 1 soil inspection pit were assessed in accordance with MAFF's revised guidelines and criteria for grading the quality of agricultural land (MAFF, 1988). These guidelines provide a framework for classifying land according to the extent to which its physical or chemical characteristics impose long term limitations on its use for agriculture.

1.3 The work was conducted by members of the Resource Planning Team in the Guildford Statutory Group of ADAS.

1.4 At the time of the survey the land use on the site was permanent grassland. An area in the north of the site is occupied by a caravan park, and is thus classed as being in Non-agricultural use.

1.5 The distribution of grades and subgrades is shown on the attached ALC map and the areas are given in the table below. The map has been drawn at a scale of 1:5,000. It is accurate at this scale, but any enlargement would be misleading. This map supersedes any previous survey information for this site.

Table 1 : Distribution of Grades and Subgrades

<u>Grade</u>	<u>Area (ha)</u>	<u>% of Site</u>	<u>% of Agricultural Area</u>
4	3.5	79.5	100% (3.5 ha)
Non-Agricultural	0.8	18.2	
Woodland	<u>0.1</u>	<u>2.3</u>	
Total area surveyed	4.4	100	

1.6 Appendix 1 gives a general description of the grades, subgrades and land use categories identified in the survey. The main classes are described in terms of the type of limitation that can occur, the typical cropping range and the expected level and consistency of yield.

1.7 All of the agricultural land surveyed has been classified as Grade 4, poor quality land, with soil wetness as the principal limitation. Heavy clay loam topsoils are underlain by poorly structured clay subsoils at shallow depths, which severely impair drainage. The interaction between these soil conditions and the local climatic regime means that these soils can be graded no higher than Grade 4.

2.0 Climate

2.1 The climatic criteria are considered first when classifying land as climate can be overriding in the sense that severe limitations will restrict land to low grades irrespective of favourable site or soil conditions.

2.2 The main parameters used in the assessment of the overall climatic limitation are average annual rainfall, as a measure of overall wetness, and accumulated temperature, as a measure of the relative warmth of a locality.

2.3 A detailed assessment of the prevailing climate was made by interpolation from a 5km gridpoint dataset (Met. Office, 1989). The details are given in the table below and these show that there is no overall climatic limitation affecting the site. However, climatic factors do interact with soil factors to influence soil wetness and soil droughtiness. At this locality, the field capacity days and average annual rainfall are relatively high thus increasing the likelihood of soil wetness.

2.4 No local climatic factors such as exposure or frost risk affect the site.

Table 2 : Climatic Interpolation

Grid Reference :	TQ 123 133
Altitude (m) :	50
Accumulated Temperature (days) :	1483
Average Annual Rainfall (mm) :	879
Field Capacity (days) :	183
Moisture Deficit, Wheat (mm) :	104
Moisture Deficit, Potatoes (mm) :	98
Overall Climatic Grade :	1

3.0 Relief

3.1 The survey area is mostly flat and lies at approximately 50m AOD. Adjacent to the stream the land gently dips to approximately 47m AOD. Nowhere on the site does gradient or relief impose any limitation to the land quality.

4.0 Geology and Soil

4.1 British Geological Sheet 318/333, Brighton and Worthing (1978) shows the northern end of the site to be underlain by Folkestone Beds and the remaining area by Gault Clay.

4.2 There are two soil types for the site, as shown on the Soil Survey map of South East England (SSEW, 1983, 1:250,000). In the north of the site, the soils comprise the Fyfield Association. These soils are described as 'well drained coarse and fine loamy soils over interbedded sands and sandstones' (SSEW, 1983). The predominant soil type on the site is shown to be the Wickham 2 Association. These soils are described as 'slowly permeable, seasonally waterlogged fine loamy over clayey and fine silty over clayey soils' (SSEW, 1983). Detailed field examination confirms drainage characteristics similar to those of the Wickham 2 Association.

5.0 Agricultural Land Classification

5.1 Table 1 provides the details of the area measurements for each grade and the distribution of each grade is shown on the attached ALC map.

5.2 The location of the soil observation points are shown on the attached sample point map.

Grade 4

5.3 All of the agricultural land surveyed has been classified as Grade 4. This poor quality land is restricted by a severe soil wetness and workability limitation. Topsoils comprise heavy clay loams and, occasionally, heavy silty clay loams. These are underlain by clay upper and lower subsoils which extend to depth. Both the upper and lower subsoils of these profiles are gleyed and poorly structured (i.e., profiles are gleyed and slowly permeable from approximately 25cm depth. These profiles are typified by Pit 1, which was found to have moderately developed coarse angular-blocky subsoil structures. Given the relatively wet climate at this locality, the slowly permeable characteristics of the subsoils severely impede drainage such that Wetness Class IV is appropriate. The interaction between topsoil textures, soil drainage characteristics and the local climatic regime means that this land can be classified as no higher than Grade 4. This soil wetness adversely affects seed germination and survival, and inhibits the development of a good root system. This limits the crops which can tolerate such conditions. In addition, severe restrictions are imposed on cultivations, grazing by livestock and trafficking by machinery.

Non-Agricultural

5.4 The Non-Agricultural land shown on the map is occupied by a caravan park.

Woodland

5.5 The Woodland marked on the map consists of mature deciduous trees.

APPENDIX I

DESCRIPTION OF THE GRADES AND SUB-GRADES

Grade 1 : Excellent Quality Agricultural Land

Land with no or very minor limitations to agricultural use. A very wide range of agricultural and horticultural crops can be grown and commonly includes top fruit, soft, fruit, salad crops and winter harvested vegetables. Yields are high and less variable than on land of lower quality.

Grade 2 : Very Good Quality Agricultural Land

Land with minor limitations which affect crop yield, cultivations or harvesting. A wide range of agricultural or horticultural crops can usually be grown but on some land on the grade there may be reduced flexibility due to difficulties with the production of the more demanding crops such as winter harvested vegetables and arable root crops. The level of yield is generally high but may be lower or more variable than Grade 1.

Grade 3 : Good To Moderate Quality Agricultural Land

Land with moderate limitations which affect the choice of crops, timing and type of cultivation, harvesting or the level of yield. When more demanding crops are grown yields are generally lower or more variable than on land in grades 1 and 2.

Sub-grade 3A : Good Quality Agricultural Land

Land capable of consistently producing moderate to high yields of a narrow range of arable crops, especially cereals, or moderate yields of a wide range of crops including cereals, grass, oilseed rape, potatoes, sugar beet and the less demanding horticultural crops.

Sub-grade 3B : Moderate Quality Agricultural Land

Land capable of producing moderate yields of a narrow range of crops, principally cereals and grass, or lower yields of a wider range of crops or high yields of grass which can be grazed or harvested over most of the year.

Grade 4 : Poor Quality Agricultural Land

Land with severe limitations which significantly restrict the range of crops and/or the level of yields. It is mainly suited to grass with occasional arable crops (eg. cereals and forage crops) the yields of which are variable. In moist climates, yields of grass may be moderate to high but there may be difficulties in utilisation. the grade also includes very droughty arable land.

Grade 5 : Very Poor Quality Agricultural Land

Land with very severe limitations which restrict use to permanent pasture or rough grazing, except for occasional pioneer forage crops.

Urban

Built-up or 'hard' uses with relatively little potential for a return to agriculture : housing, industry, commerce, education, transport, religious buildings, cemeteries. Also, hard-surfaced sports facilities, permanent caravan sites and vacant land; all types of derelict land, including mineral workings which are only likely to be reclaimed using derelict land grants.

Non-agricultural

'Soft' uses where most of the land could be returned relatively easily to agriculture, including : private parkland, public open spaces, sports fields, allotments and soft-surfaced areas on airports/airfields. Also active mineral workings and refuse tips where restoration conditions to 'soft' after-uses may apply.

Woodland

Includes commercial and non-commercial woodland.

Agricultural Buildings

Includes the normal range of agricultural buildings as well as other relatively permanent structures such as glasshouses. Temporary structures (eg. polythene tunnels erected for lambing) may be ignored.

Open Water

Includes lakes, ponds and rivers as map scale permits.

Land Not Surveyed

Agricultural land which has not been surveyed.

Where the land use includes more than one of the above, eg. buildings in large grounds, and where map scale permits, the cover types may be shown separately. Otherwise, the most extensive cover type will be shown.

APPENDIX II

REFERENCES

- * British Geological Survey (1978), Sheet No. 318/333, Brighton and Worthing, 1:50,000.
- * MAFF (1988), Agricultural Land Classification of England And Wales : Revised guidelines and criteria for grading the quality of agricultural land.
- * Meteorological Office (1989), Climatological Data for Agricultural Land Classification.
- * Soil Survey of England and Wales (1983), Sheet 6, Soils of South East England, 1:250,000 and accompanying legend.

APPENDIX III

DEFINITION OF SOIL WETNESS CLASSES

Wetness Class I

The soil profile is not wet within 70cm depth for more than 30 days in most years.

Wetness Class II

The soil profile is wet within 70cm depth for 31-90 days in most years or, if there is no slowly permeable layer within 80cm depth, it is wet within 70cm for more than 90 days, but not wet within 40cm depth for more than 30 days in most years.

Wetness Class III

The soil profile is wet within 70cm depth for 91-180 days in most years or, if there is no slowly permeable layer within 80cm depth, it is wet within 70cm for more than 180 days, but only wet within 40cm depth for 31-90 days in most years.

Wetness Class IV

The soil profile is wet within 70cm depth for more than 180 days but not wet within 40cm depth for more than 210 days in most years or, if there is no slowly permeable layer within 80cm depth, it is wet within 40cm depth for 91-210 days in most years.

Wetness Class V

The soil profile is wet within 40cm depth for 211-335 days in most years.

Wetness Class VI

The soil profile is wet within 40cm depth for more than 335 days in most years.

(The number of days is not necessarily a continuous period. 'In most years' is defined as more than 10 out of 20 years.)

APPENDIX IV

SOIL PIT AND SOIL BORING DESCRIPTIONS

- Contents :
- * Soil Abbreviations : Explanatory Note
 - * Soil Pit Descriptions
 - * Database Printout : Boring Level Information
 - * Database Printout : Horizon Level Information

SOIL PROFILE DESCRIPTIONS : EXPLANATORY NOTE

Soil pit and auger boring information collected during ALC fieldwork is held on a database. This has commonly used notations and abbreviations as set out below.

Boring Header Information

1. **GRID REF** : national grid square and 8 figure grid reference.

2. **USE** : Land use at the time of survey. The following abbreviations are used.

ARA : Arable **WHT** : Wheat **BAR** : Barley **CER** : Cereals **OAT** : Oats **MZE** : Maize **OSR** : Oilseed rape
BEN : Field Beans **BRA** : Brassicae **POT** : Potatoes **SBT** : Sugar Beet **FCD** : Fodder Crops **LIN** : Linseed
FRT : Soft and Top Fruit **HRT** : Horticultural Crops **PGR** : Permanent Pasture **LEY** : Ley Grass **RGR** : Rough Grazing
SCR : Scrub **CFW** : Coniferous Woodland **DCW** : Deciduous Woodland **HTH** : Heathland **BOG** : Bog or Marsh
FLW : Fallow **PLO** : Ploughed **SAS** : Set aside **OTH** : Other

3. **GRDNT** : Gradient as measured by a hand-held optical clinometer.

4. **GLEYSPL** : Depth in cm to gleying or slowly permeable layers.

5. **AP (WHEAT/POTS)** : Crop-adjusted available water capacity.

6. **MB (WHEAT/POTS)** : Moisture Balance.

7. **DRT** : Best grade according to soil droughtiness.

8. If any of the following factors are considered significant, an entry of 'Y' will be entered in the relevant column.

MREL : Microrelief limitation **FLOOD** : Flood risk **EROSN** : Soil erosion risk **EXP** : Exposure limitation **FROST** : Frost
DIST : Disturbed land **CHEM** : Chemical limitation

9. **LIMIT** : The main limitation to land quality. The following abbreviations are used.

OC : Overall Climate **AE** : Aspect **EX** : Exposure **FR** : Frost Risk **GR** : Gradient **MR** : Microrelief
FL : Flood Risk **TX** : Topsoil Texture **DP** : Soil Depth **CH** : Chemical **WE** : Wetness **WK** : Workability
DR : Drought **ER** : Soil Erosion Risk **WD** : Combined Soil Wetness/Droughtiness **ST** : Topsoil Stoniness

Soil Pits and Auger Borings

1. **TEXTURE** : soil texture classes are denoted by the following abbreviations.

S : Sand **LS** : Loamy Sand **SL** : Sandy Loam **SZL** : Sandy Silt Loam **CL** : Clay Loam **ZCL** : Silty Clay Loam
SCL : Sandy Clay Loam **C** : Clay **SC** : Sandy Clay **ZC** : Silty Clay **OL** : Organic Loam **P** : Peat **SP** : Sandy Peat
LP : Loamy Peat **PL** : Peaty Loam **PS** : Peaty Sand **MZ** : Marine Light Silts

For the sand, loamy sand, sandy loam and sandy silt loam classes, the predominant size of sand fraction will be indicated by the use of prefixes.

F : Fine (more than 66% of the sand less than 0.2mm)

M : Medium (less than 66% fine sand and less than 33% coarse sand)

C : Coarse (more than 33% of the sand larger than 0.6mm)

The clay loam and silty clay loam classes will be sub-divided according to the clay content.

M : Medium (< 27% clay) **H** : Heavy (27-35% clay)

2. **MOTTLE COL** : Mottle colour

3. **MOTTLE ABUN** : Mottle abundance, expressed as a percentage of the matrix or surface described.

F : few <2% C : common 2-20% M : many 20-40 VM : very many 40% +

4. **MOTTLE CONT** : Mottle contrast

F : faint - indistinct mottles, evident only on close inspection D : distinct - mottles are readily seen

P : prominent - mottling is conspicuous and one of the outstanding features of the horizon

5. **PED. COL** : Ped face colour

6. **STONE LITH** : One of the following is used.

HR : all hard rocks and stones MSST : soft, medium or coarse grained sandstone

SI : soft weathered igneous or metamorphic SLST : soft oolitic or dolimitic limestone

FSST : soft, fine grained sandstone ZR : soft, argillaceous, or silty rocks CH : chalk

GH : gravel with non-porous (hard) stones GS : gravel with porous (soft) stones

Stone contents (> 2cm, > 6cm and total) are given in percentages (by volume).

7. **STRUCT** : the degree of development, size and shape of soil peds are described using the following notation:

- degree of development WK : weakly developed MD : moderately developed ST : strongly developed

- ped size F : fine M : medium C : coarse VC : very coarse

- ped shape S : single grain M : massive GR : granular AB : angular blocky SAB : sub-angular blocky PR : prismatic
PL : platy

8. **CONSIST** : Soil consistence is described using the following notation:

L : loose VF : very friable FR : friable FM : firm VM : very firm EM : extremely firm EH : extremely hard

9. **SUBS STR** : Subsoil structural condition recorded for the purpose of calculating profile droughtiness.

G : good M : moderate P : poor

10. **POR** : Soil porosity. If a soil horizon has less than 0.5% biopores > 0.5 mm, a 'Y' will appear in this column.

11. **IMP** : If the profile is impenetrable a 'Y' will appear in this column at the appropriate horizon.

12. **SPL** : Slowly permeable layer. If the soil horizon is slowly permeable a 'Y' will appear in this column.

13. **CALC** : If the soil horizon is calcareous, a 'Y' will appear in this column.

14. Other notations

APW : available water capacity (in mm) adjusted for wheat

APP : available water capacity (in mm) adjusted for potatoes

MBW : moisture balance, wheat

MBP : moisture balance, potatoes

SOIL PIT DESCRIPTION

Site Name : WSUSSEX MINS SITE 15 Pit Number : 1P

Grid Reference: TQ12411319 Average Annual Rainfall : 879 mm
 Accumulated Temperature : 1483 degree days
 Field Capacity Level : 183 days
 Land Use : Permanent Grass
 Slope and Aspect : degrees

HORIZON	TEXTURE	COLOUR	STONES >2	TOT.STONE	MOTTLES	STRUCTURE
0- 24	HCL	25Y 32 00	0	0	F	MDCSAB
24- 49	C	25Y 52 00	0	0	M	MDCAB
49-120	C	25Y 51 00	0	5	M	MDCAB

Wetness Grade : 4 Wetness Class : IV
 Gleying : 024 cm
 SPL : 024 cm

Drought Grade : 2 APW : 124mm MBW : 20 mm
 APP : 102mm MBP : 6 mm

FINAL ALC GRADE : 4
 MAIN LIMITATION : Wetness

SAMPLE NO.	GRID REF	ASPECT USE	--WETNESS--		-WHEAT-		-POTS-		M.REL		EROSN	FROST		CHEM	ALC	COMMENTS
			GRDNT	GLEYSPL	CLASS	GRADE	AP	MB	AP	MB	DRT	FLOOD	EXP	DIST	LIMIT	
1P	TQ12411319	PGR	024	024	4	4	124	20	102	6	2				WE 4	PIT DUG TO 120
2	TQ12301330	PGR	025	025	4	4	126	22	103	7	2				WE 4	GLEYSPL 25
3	TQ12391330	PGR	020		2	3A	151	47	111	15	1				WE 3A	EDGE OF SITE
4	TQ12401320	PGR	025	025	4	4	124	20	114	18	2				WE 4	GLEYSPL 25
5	TQ12501310	PGR	0	028	4	4	127	23	104	8	2				WE 4	GLEYSPL 28
6	TQ12391327	PGR	030	030	4	4	129	25	106	10	2				WE 4	GLEYSPL 30

SAMPLE	DEPTH	TEXTURE	COLOUR	----MOTTLES-----			PED		----STONES-----			STRUCT/ CONSIST	SUBS					
				COL	ABUN	CONT	COL.	GLE	>2	>6	LITH		TOT	STR	POR	IMP	SPL	CALC
1P	0-24	hc1	25Y 32 00	75YR46	00	F			0	0	0	MDCSAB	FM					
	24-49	c	25Y 52 00	10YR56	00	M		Y	0	0	0	MDCAB	FM P	Y		Y		
	49-120	c	25Y 51 00	10YR58	00	M	10YR52	00	Y	0	0	HR	5	MDCAB	VM P	Y	Y	
2	0-25	hc1	25Y 42 00						0	0	0							
	25-75	c	10YR62	53	10YR58	52	M	Y	0	0	HR	1		P	Y	Y		
	75-95	c	10YR53	00	10YR56	00	M	Y	0	0	HR	1		P	Y	Y		
	95-120	c	25Y 42 00	10YR56	00	C		Y	0	0	CH	5		P	Y	Y		
3	0-20	hc1	10YR32	00					0	0	0							
	20-120	sc1	10YR62	00	10YR68	00	C	Y	0	0	0			M				
4	0-25	hzc1	10YR32	00					0	0	0							
	25-100	zc	10YR51	00	10YR68	00	C	Y	0	0	CH	5		M			Y	
5	0-28	hc1	25Y 42 00	10YR58	00	C		Y	0	0	0							
	28-80	c	10YR62	53	10YR56	00	M	00MNO0	00	Y	0	0	HR	2		P	Y	Y
	80-120	c	25Y 51 00	10YR58	00	M		Y	0	0	CH	10		P	Y	Y	Y	
6	0-30	hc1	10YR42	00					0	0	0							
	30-65	c	10YR53	00	10YR56	52	M	Y	0	0	HR	1		P	Y	Y	Y	
	65-120	c	25Y 42 00	10YR58	00	C		Y	0	0	CH	5		P	Y	Y	Y	