

A1
ARUN DISTRICT
LOCAL PLAN : SITE 29
LAND EAST OF
YAPTON ROAD, CLIMPING
AGRICULTURAL LAND CLASSIFICATION
ALC MAP & REPORT
MARCH 1994

**ARUN DISTRICT LOCAL PLAN : SITE 29 LAND EAST OF
YAPTON ROAD, CLIMPING
AGRICULTURAL LAND CLASSIFICATION REPORT**

1.0 Summary

1.1 ADAS was commissioned by MAFF's Land Use Planning Unit to provide information on land quality of a number of sites in the Arun District of West Sussex. The work forms part of MAFF's statutory input to the preparation of the Arun District Local Plan.

1.2 Approximately 7 hectares of land relating to site 29 at the disused aerodrome near the village of Yapton in West Sussex was surveyed in March 1994. The survey was undertaken at a detailed level of approximately one boring per hectare. A total of 6 soil auger borings and one soil inspection pit were assessed in accordance with MAFF's revised guidelines and criteria for grading the quality of agricultural land (MAFF, 1988). These guidelines provide a framework for classifying land according to the extent to which its physical or chemical characteristics impose long term limitations on its use for agriculture.

1.3 The work was conducted by members of the Resource Planning Team in the Guildford Statutory Group of ADAS.

1.4 At the time of the survey the land was under a cereal crop.

1.5 The distribution of grades and subgrades is shown on the attached ALC map and the areas are given in the table below. The map has been drawn at a scale of 1:5,000. It is accurate at this scale, but any enlargement would be misleading. This map supersedes any previous survey information for this site.

Table 1 : Distribution of Grades and Subgrades

<u>Grade</u>	<u>Area (ha)</u>	<u>% of Site</u>
2	1.8	25
3a	5.4	75
Total Area of Site	7.2	100%

1.6 Appendix 1 gives a general description of the grades, subgrades and land use categories identified in the survey. The main classes are described in terms of the type of limitation that can occur, the typical cropping range and the expected level and consistency of yield.

1.7 The site is classified as grades 2 and 3a with soil wetness and droughtiness being the main limitations. The majority of the land is grade 2 and comprises fine loamy soils passing to slowly permeable clay at depth. As a result water drainage through these soils is slightly impaired and land is classified as grade 2 due to a slight soil wetness limitation. Together with this, the comparatively dry nature of the local climate interacts with these soils to produce a slight restriction in available water reserves for plant growth. Therefore land is also limited to grade 2 due to a slight droughtiness limitation. Subgrade 3a land consists of similar soils but with slowly permeable clay at a shallower depth in the profile. Consequently drainage is more severely impaired than that of grade 2 land and this land is classified as subgrade 3a due to a moderate wetness limitation.

2.0 Climate

2.1 The climatic criteria are considered first when classifying land as climate can be overriding in the sense that severe limitations will restrict land to low grades irrespective of favourable site or soil conditions.

2.2 The main parameters used in the assessment of an overall climatic limitation are annual average rainfall, as a measure of overall wetness, and accumulated temperature, as a measure of the relative warmth of a locality.

2.3 A detailed assessment of the prevailing climate was made by interpolation from a 5km gridpoint dataset (Met. Office 1989). The details are given in the table below and these show that there is no overall climatic limitation affecting the site.

2.4 No local climatic factors such as exposure or frost risk affect the site. However, it should be noted that climatic characteristics can interact with soil properties to influence soil wetness and droughtiness.

Table 2 : Climatic Interpolation

Grid Reference :	SU 992 024
Altitude (m) :	5
Accumulated Temperature (days) :	1543
Average Annual Rainfall (mm) :	742
Field Capacity (days) :	152
Moisture Deficit, Wheat (mm) :	121
Moisture Deficit, Potatoes (mm) :	118
Overall Climatic Grade :	1

3.0 Relief

3.1 The site is flat and lies at an altitude of approximately 5 metres AOD. Nowhere on the site do relief or gradient affect agricultural land quality.

4.0 Geology and Soil

4.1 The published geological sheet for the site, Sheet 332 (BGS, 1975) shows the underlying geology to be brickearth over Upper Chalk.

4.2 The published soils information for the area, Sheet SU90 (SSGB, 1967) shows the majority of the site to comprise soils of the Park Gate Series which is described as "Deep stoneless silty soils variably affected by groundwater" (SSEW, 1983). To the north east of the site is mapped soils of the Hook Series, described as "Deep stoneless well drained silty soils and similar soils affected by groundwater; over gravel locally. Usually flat land" (SSEW, 1983). A detailed inspection of soils on the site found fine loamy soils over slowly permeable clay with associated wetness problems.

5.0 Agricultural Land Classification

5.1 Table 1 provides the details of the area measurements for each grade and the distribution of each grade is shown on the attached ALC map.

5.2 The locations of the soil observation points are shown on the attached sample point map.

Grade 2

5.3 Very good quality land covers the majority of the site. Typical soil profiles comprise medium clay loam, occasionally medium silty clay loam topsoils containing 0-2% total flints over very slightly stony upper subsoils of heavy clay loam, occasionally medium clay loam. This passes to poorly structured slowly permeable clay towards the bottom of the profile. Soil Pit 1 is typical of these soils. Wetness imperfections in the form of gleying were observed from 29-75 cm caused by slowly permeable horizons of clay beginning from 75 cm depth. Consequently, soils are assigned to a wetness class of II, reflecting their drainage status, which together with a medium topsoil texture at this location gives a grade of 2, the land being limited by slight soil wetness.

5.4 In addition to this limitation, land is also limited to grade 2 due to slight soil droughtiness. The interaction of soil properties with climatic characteristics results in a slight restriction of profile available water reserves. Nevertheless such land is suitable for a wide range of agricultural and horticultural uses.

Subgrade 3a

5.5 Subgrade 3a, good quality agricultural land comprises soils similar to that of grade 2 land. Topsoils consist of medium clay loam or medium silty clay loam with 0-3% total flints over thin upper subsoils of stoneless heavy clay loam, passing to poorly structured slowly permeable clay. Soils are imperfectly drained (wetness class III), being gleyed from 30-35 cm depth due to slowly permeable horizons of clay beginning from 45-60 cm depth. Land is classified as subgrade 3a due to a moderate wetness limitation. The grade reflects the increasing wetness limitation making land slightly less flexible than that graded 2.

ADAS REFERENCE : 4202/065/94
MAFF REFERENCE : EL 42/460

Resource Planning Team
Guildford Statutory Group
ADAS Reading

APPENDIX I

DESCRIPTION OF THE GRADES AND SUB-GRADES

Grade 1 : Excellent Quality Agricultural Land

Land with no or very minor limitations to agricultural use. A very wide range of agricultural and horticultural crops can be grown and commonly includes top fruit, soft, fruit, salad crops and winter harvested vegetables. Yields are high and less variable than on land of lower quality.

Grade 2 : Very Good Quality Agricultural Land

Land with minor limitations which affect crop yield, cultivations or harvesting. A wide range of agricultural or horticultural crops can usually be grown but on some land on the grade there may be reduced flexibility due to difficulties with the production of the more demanding crops such as winter harvested vegetables and arable root crops. The level of yield is generally high but may be lower or more variable than Grade 1.

Grade 3 : Good To Moderate Quality Agricultural Land

Land with moderate limitations which affect the choice of crops, timing and type of cultivation, harvesting or the level of yield. When more demanding crops are grown yields are generally lower or more variable than on land in grades 1 and 2.

Sub-grade 3A : Good Quality Agricultural Land

Land capable of consistently producing moderate to high yields of a narrow range of arable crops, especially cereals, or moderate yields of a wide range of crops including cereals, grass, oilseed rape, potatoes, sugar beet and the less demanding horticultural crops.

Sub-grade 3B : Moderate Quality Agricultural Land

Land capable of producing moderate yields of a narrow range of crops, principally cereals and grass, or lower yields of a wider range of crops or high yields of grass which can be grazed or harvested over most of the year.

Grade 4 : Poor Quality Agricultural Land

Land with severe limitations which significantly restrict the range of crops and/or the level of yields. It is mainly suited to grass with occasional arable crops (eg. cereals and forage crops) the yields of which are variable. In moist climates, yields of grass may be moderate to high but there may be difficulties in utilisation. the grade also includes very droughty arable land.

Grade 5 : Very Poor Quality Agricultural Land

Land with very severe limitations which restrict use to permanent pasture or rough grazing, except for occasional pioneer forage crops.

Urban

Built-up or 'hard' uses with relatively little potential for a return to agriculture : housing, industry, commerce, education, transport, religious buildings, cemeteries. Also, hard-surfaced sports facilities, permanent caravan sites and vacant land; all types of derelict land, including mineral workings which are only likely to be reclaimed using derelict land grants.

Non-agricultural

'Soft' uses where most of the land could be returned relatively easily to agriculture, including : private parkland, public open spaces, sports fields, allotments and soft-surfaced areas on airports/airfields. Also active mineral workings and refuse tips where restoration conditions to 'soft' after-uses may apply.

Woodland

Includes commercial and non-commercial woodland.

Agricultural Buildings

Includes the normal range of agricultural buildings as well as other relatively permanent structures such as glasshouses. Temporary structures (eg. polythene tunnels erected for lambing) may be ignored.

Open Water

Includes lakes, ponds and rivers as map scale permits.

Land Not Surveyed

Agricultural land which has not been surveyed.

Where the land use includes more than one of the above, eg. buildings in large grounds, and where map scale permits, the cover types may be shown separately. Otherwise, the most extensive cover type will be shown.

APPENDIX II

REFERENCES

- * BRITISH GEOLOGICAL SURVEY (1975), Sheet No.332, Bognor Regis, 1:50,000 scale.
- * MAFF (1988), Agricultural Land Classification of England And Wales : Revised guidelines and criteria for grading the quality of agricultural land.
- * METEOROLOGICAL OFFICE (1989), Climatological Data for Agricultural Land Classification.
- * SOIL SURVEY OF ENGLAND AND WALES (1983), Sheet No.6, "Soils of South East England", 1:250,000 scale and accompanying legend.
- * SOIL SURVEY OF GREAT BRITAIN (1967), Bulletin No.3, Soils of the West Sussex Coastal Plain.

APPENDIX III

DEFINITION OF SOIL WETNESS CLASSES

Wetness Class I

The soil profile is not wet within 70cm depth for more than 30 days in most years.

Wetness Class II

The soil profile is wet within 70cm depth for 31-90 days in most years or, if there is no slowly permeable layer within 80cm depth, it is wet within 70cm for more than 90 days, but not wet within 40cm depth for more than 30 days in most years.

Wetness Class III

The soil profile is wet within 70cm depth for 91-180 days in most years or, if there is no slowly permeable layer within 80cm depth, it is wet within 70cm for more than 180 days, but only wet within 40cm depth for 31-90 days in most years.

Wetness Class IV

The soil profile is wet within 70cm depth for more than 180 days but not wet within 40cm depth for more than 210 days in most years or, if there is no slowly permeable layer within 80cm depth, it is wet within 40cm depth for 91-210 days in most years.

Wetness Class V

The soil profile is wet within 40cm depth for 211-335 days in most years.

Wetness Class VI

The soil profile is wet within 40cm depth for more than 335 days in most years.

(The number of days is not necessarily a continuous period. 'In most years' is defined as more than 10 out of 20 years.)

APPENDIX IV

SOIL PIT AND SOIL BORING DESCRIPTIONS

- Contents :
- * Soil Abbreviations : Explanatory Note
 - * Soil Pit Descriptions
 - * Database Printout : Boring Level Information
 - * Database Printout : Horizon Level Information

SOIL PROFILE DESCRIPTIONS : EXPLANATORY NOTE

Soil pit and auger boring information collected during ALC fieldwork is held on a database. This has commonly used notations and abbreviations as set out below.

Boring Header Information

1. **GRID REF** : national grid square and 8 figure grid reference.

2. **USE** : Land use at the time of survey. The following abbreviations are used.

ARA : Arable **WHT** : Wheat **BAR** : Barley **CER** : Cereals **OAT** : Oats **MZE** : Maize **OSR** : Oilseed rape
BEN : Field Beans **BRA** : Brassicae **POT** : Potatoes **SBT** : Sugar Beet **FCD** : Fodder Crops **LIN** : Linseed
FRT : Soft and Top Fruit **HRT** : Horticultural Crops **PGR** : Permanent Pasture **LEY** : Ley Grass **RGR** : Rough Grazing
SCR : Scrub **CFW** : Coniferous Woodland **DCW** : Deciduous Woodland **HTH** : Heathland **BOG** : Bog or Marsh
FLW : Fallow **PLO** : Ploughed **SAS** : Set aside **OTH** : Other

3. **GRDNT** : Gradient as measured by a hand-held optical clinometer.

4. **GLEY/SPL** : Depth in cm to gleying/slight gleying or slowly permeable layers.

5. **AP (WHEAT/POTS)** : Crop-adjusted available water capacity.

6. **MB (WHEAT/POTS)** : Moisture Balance.

7. **DRT** : Best grade according to soil droughtiness.

8. If any of the following factors are considered significant, an entry of 'Y' will be entered in the relevant column.

MREL : Microrelief limitation **FLOOD** : Flood risk **EROSN** : Soil erosion risk **EXP** : Exposure limitation **FROST** : Frost
DIST : Disturbed land **CHEM** : Chemical limitation

9. **LIMIT** : The main limitation to land quality. The following abbreviations are used.

OC : Overall Climate **AE** : Aspect **EX** : Exposure **FR** : Frost Risk **GR** : Gradient **MR** : Microrelief
FL : Flood Risk **TX** : Topsoil Texture **DP** : Soil Depth **CH** : Chemical **WE** : Wetness **WK** : Workability
DR : Drought **ER** : Soil Erosion Risk **WD** : Combined Soil Wetness/Droughtiness **ST** : Topsoil Stoniness

Soil Pits and Auger Borings

1. **TEXTURE** : soil texture classes are denoted by the following abbreviations.

S : Sand **LS** : Loamy Sand **SL** : Sandy Loam **SZL** : Sandy Silt Loam **CL** : Clay Loam **ZCL** : Silty Clay Loam
SCL : Sandy Clay Loam **C** : Clay **SC** : Sandy Clay **ZC** : Silty Clay **OL** : Organic Loam **P** : Peat **SP** : Sandy Peat
LP : Loamy Peat **PL** : Peaty Loam **PS** : Peaty Sand **MZ** : Marine Light Silts

For the sand, loamy sand, sandy loam and sandy silt loam classes, the predominant size of sand fraction will be indicated by the use of prefixes.

F : Fine (more than 66% of the sand less than 0.2mm)

M : Medium (less than 66% fine sand and less than 33% coarse sand)

C : Coarse (more than 33% of the sand larger than 0.6mm)

The clay loam and silty clay loam classes will be sub-divided according to the clay content.

M : Medium (< 27% clay) **H** : Heavy (27-35% clay)

2. **MOTTLE COL** : Mottle colour

3. **MOTTLE ABUN** : Mottle abundance, expressed as a percentage of the matrix or surface described.

F : few <2% C : common 2-20% M : many 20-40 VM : very many 40%+

4. **MOTTLE CONT** : Mottle contrast

F : faint - indistinct mottles, evident only on close inspection D : distinct - mottles are readily seen

P : prominent - mottling is conspicuous and one of the outstanding features of the horizon

5. **PED. COL** : Ped face colour

6. **STONE LITH** : One of the following is used.

HR : all hard rocks and stones MSST : soft, medium or coarse grained sandstone

SI : soft weathered igneous or metamorphic SLST : soft oolitic or dolimitic limestone

FSST : soft, fine grained sandstone ZR : soft, argillaceous, or silty rocks CH : chalk

GH : gravel with non-porous (hard) stones GS : gravel with porous (soft) stones

Stone contents (> 2cm, > 6cm and total) are given in percentages (by volume).

7. **STRUCT** : the degree of development, size and shape of soil peds are described using the following notation:

- degree of development WK : weakly developed MD : moderately developed ST : strongly developed

- ped size F : fine M : medium C : coarse VC : very coarse

- ped shape S : single grain M : massive GR : granular AB : angular blocky SAB : sub-angular blocky PR : prismatic
PL : platy

8. **CONSIST** : Soil consistence is described using the following notation:

L : loose VF : very friable FR : friable FM : firm VM : very firm EM : extremely firm EH : extremely hard

9. **SUBS STR** : Subsoil structural condition recorded for the purpose of calculating profile droughtiness.

G : good M : moderate P : poor

10. **POR** : Soil porosity. If a soil horizon has less than 0.5% biopores > 0.5 mm, a 'Y' will appear in this column.

11. **IMP** : If the profile is impenetrable a 'Y' will appear in this column at the appropriate horizon.

12. **SPL** : Slowly permeable layer. If the soil horizon is slowly permeable a 'Y' will appear in this column.

13. **CALC** : If the soil horizon is calcareous, a 'Y' will appear in this column.

14. Other notations

APW : available water capacity (in mm) adjusted for wheat

APP : available water capacity (in mm) adjusted for potatoes

MBW : moisture balance, wheat

MBP : moisture balance, potatoes

SOIL PIT DESCRIPTION

Site Name : ARUN LOCAL PLAN SITE 29 Pit Number : 1P

Grid Reference: SU99300230 Average Annual Rainfall : 742 mm
 Accumulated Temperature : 1543 degree days
 Field Capacity Level : 152 days
 Land Use : Cereals
 Slope and Aspect : degrees

HORIZON	TEXTURE	COLOUR	STONES >2	TOT.STONE	MOTTLES	STRUCTURE
0- 32	MCL	10YR42 00	0	0		
32- 57	MCL	25Y 54 00	0	0		MDCSAB
57- 80	HCL	25Y 53 00	0	0	C	MDCSAB
80-100	C	25Y 64 00	0	0	M	WKMPR
100-120	HCL	25Y 64 00	0	0	M	

Wetness Grade : 2 Wetness Class : II
 Gleying : 057 cm
 SPL : 080 cm

Drought Grade : 2 APW : 150mm MBW : 29 mm
 APP : 118mm MBP : 0 mm

FINAL ALC GRADE : 2
 MAIN LIMITATION : Wetness

SAMPLE NO.	GRID REF	ASPECT USE	--WETNESS--		-WHEAT-		-POTS-		M. REL		EROSN EXP	FROST DIST	CHEM LIMIT	ALC	COMMENTS
			GRDNT	GLEYSPL	CLASS	GRADE	AP	MB	AP	MB					
1	SU99100250	CER	070	080	2	2	143	22	117	-1	2		WE	2	WEDR
1P	SU99300230	CER	057	080	2	2	150	29	118	0	2		WE	2	
2	SU99200240	CER	055	090	1	1	147	26	118	0	2		DR	2	
3	SU99300240	CER	030	060	3	3A	136	15	113	-5	2		WE	3A	
4	SU99300230	CER	075	075	2	2	141	20	117	-1	2		WE	2	WEDR
5	SU99400230	CER	035	045	3	3A	150	29	112	-6	2		WE	3A	SLI GLEY 29
6	SU99500230	CER	029	078	2	2	150	29	119	1	2		WE	2	WEDR

SAMPLE	DEPTH	TEXTURE	COLOUR	----MOTTLES----- PED			----STONES-----			STRUCT/	SUBS	SPL	CALC	
				COL	ABUN	CONT	COL.	GLE	>2					>6
1	0-25	mc1	10YR43 00						0	0	HR	1		
	25-50	mc1	10YR53 00						0	0		0	M	
	50-70	hc1	10YR53 00	00MNOO	00	F			0	0		0	M	
	70-80	hc1	10YR53 00	10YR58	61	C	00MNOO	00	Y	0	0	0	M	
	80-120	c	10YR64 00	10YR68	71	M	00MNOO	00	Y	0	0	0	P	Y
1P	0-32	mc1	10YR42 00						0	0		0		
	32-57	mc1	25Y 54 00						0	0		0	MDCSAB FR M	
	57-80	hc1	25Y 53 00	10YR58	00	C	00MNOO	00	Y	0	0	0	MDCSAB FM M	
	80-100	c	25Y 64 00	10YR68	71	M	00MNOO	00	Y	0	0	0	WKMPR FM P	Y
	100-120	hc1	25Y 64 00	10YR68	71	M	00MNOO	00	Y	0	0	0	M	ASSUME M-NO SPL
2	0-35	mc1	75YR53 00						0	0	HR	1		
	35-55	mc1	10YR54 00						0	0		0	M	
	55-90	hc1	10YR53 00	10YR58	61	C	00MNOO	00	Y	0	0	0	M	
	90-120	c	10YR64 00	10YR68	71	M	00MNOO	00	Y	0	0	0	P	Y
3	0-30	mc1	10YR42 00						0	0	HR	3		
	30-60	hc1	10YR53 00	10YR58	00	C	00MNOO	00	Y	0	0	0	M	
	60-120	c	10YR64 00	10YR68	71	M	00MNOO	00	Y	0	0	0	P	Y
4	0-29	mc1	10YR42 00						0	0	HR	2		
	29-50	mc1	25Y 54 00						0	0		0	M	
	50-75	hc1	25Y 64 00	10YR56	00	F			0	0		0	M	
	75-120	c	25Y 63 64	75YR56	00	C			Y	0	0	0	P	Y
5	0-29	mzc1	10YR43 00						0	0	HR	2		
	29-35	mc1	10YR66 00	10YR56	00	C			S	0	0	0	M	SLI GLEY
	35-45	hc1	10YR53 00	75YR56	00	M			Y	0	0	0	M	
	45-70	c	10YR53 54	75YR56	00	C			Y	0	0	0	P	Y
	70-120	hc1	10YR53 54	75YR56	00	C			Y	0	0	0	M	
6	0-29	mzc1	10YR43 00						0	0	HR	2		
	29-48	mc1	10YR53 00	10YR56	00	C			Y	0	0	0	M	
	48-78	hc1	10YR53 54	10YR56	00	C			Y	0	0	HR	1	M
	78-100	c	10YR53 54	75YR56	00	C			Y	0	0	0	P	Y
	100-120	hc1	10YR53 54	10YR56	00	C			Y	0	0	HR	5	M