

**A1**  
**Maidstone Borough Local Plan**  
**Site 46 Vicarage Field,**  
**Linton Hill, Linton**  
**Agricultural Land Classification**  
**ALC Map and Report**  
**September 1994**

# AGRICULTURAL LAND CLASSIFICATION REPORT

## MAIDSTONE BOROUGH LOCAL PLAN

### SITE 46 VICARAGE FIELD, LINTON HILL, LINTON

#### 1 Summary

1.1 ADAS was commissioned by MAFF's Land Use Planning Unit to provide information on land quality for a number of sites in the Maidstone Borough of Kent. The work forms part of MAFF's statutory input to the preparation of the Maidstone Borough Local Plan.

1.2 Site 46 comprises 1.9 hectares of land west of Linton Hill and north of Wheelers Lane in the village of Linton. An Agricultural Land Classification (ALC) survey was carried out during August 1994. The survey was undertaken at a detailed level of approximately two borings per hectare of agricultural land surveyed. A total of 4 borings and one soil inspection pit were described in accordance with MAFF's revised guidelines and criteria for grading the quality of agricultural land (MAFF 1988). These guidelines provide a framework for classifying land according to the extent to which its physical or chemical characteristics impose a long term limitation on its use for agriculture.

1.3 At the time of the survey the land use on the site was pear orchards.

1.4 The site is shown on the attached ALC map which has been drawn at a scale of 1:10,000. It is accurate at this scale but any enlargement would be misleading.

1.5 Appendix I gives a general description of the grades, subgrades and land use categories identified in the survey. The main classes are described in terms of the type of limitation that can occur, the typical cropping range and the expected level and consistency of yield.

1.6 All of the agricultural land surveyed (1.9 ha) has been classified as Subgrade 3b moderate quality because of significant soil wetness and workability limitations. Profiles typically comprise non-calcareous heavy clay loams and heavy silty clay loams over similarly textured upper subsoils and clay lower subsoils. The clay is poorly structured and its slowly permeable characteristics act to significantly impair drainage, giving rise to soil wetness which may adversely affect crop growth and the utilisation of the land.

#### 2 Climate

2.1 The climatic criteria are considered first when classifying land as climate can be overriding in the sense that severe limitations will restrict land to low grades irrespective of favourable site or soil conditions.

- 2 2 The main parameters used in the assessment of an overall climatic limitation are average annual rainfall as a measure of overall wetness and accumulated temperature (degree days Jan-June) as a measure of the relative warmth of a locality
- 2 3 A detailed assessment of the prevailing climate was made by interpolation from a 5km gridpoint dataset (Met Office 1989) The details are given in the table below and these show that there is no overall climatic limitation affecting the site However in a regional context the crop adjusted soil moisture deficits are relatively high and the field capacity days are relatively low at this locality These climatic factors respectively increase the likelihood of soil droughtiness limitations and decrease the likelihood of soil wetness limitations
- 2 4 No local climatic factors such as exposure or frost risk are believed to affect the site

**Table 1 Climatic Interpolation**

Grid Reference	TQ754496
Altitude (m)	55
Accumulated Temperature (degree days Jan June)	1447
Average Annual Rainfall (mm)	681
Field Capacity (days)	138
Moisture Deficit Wheat (mm)	119
Moisture Deficit Potatoes (mm)	114
Overall Climatic Grade	1

### **3 Relief**

- 3 1 The site occupies a gentle hillside falling through gradients of 2° from approximately 57m AOD along the northern site boundary to lie at approximately 50m AOD along the southern site boundary Neither gradient or relief impose any limitation to agricultural land quality

### **4 Geology and Soil**

- 4 1 The relevant geological sheet (BGS 1976) shows the entire site to be underlain by Weald Clay
- 4 2 The published Soil Survey map (SSEW 1983) shows the Wickham 1 association at this site These soils are described as slowly permeable seasonally waterlogged fine silty over clayey fine loamy over clayey and clayey soils (SSEW 1983)
- 4 3 Detailed field examination found one soil type comprising heavy textured poorly drained soils across the entire site

## 5 **Agricultural Land Classification**

- 5 1 The distribution of Subgrade 3b land is shown on the attached ALC map
- 5 2 The location of the soil observation points are shown on the attached sample point map

### **Subgrade 3b**

- 5 3 All of the land has been classed as moderate quality because of significant soil wetness and workability limitations. Topsoils typically comprise very slightly stony non calcareous heavy clay loams and heavy silty clay loams. These are underlain by similarly textured very slightly stony moderately structured upper subsoils. At approximately 36 cm depth though occasionally deeper these profiles pass into very slightly to slightly stony clay lower subsoils. These clays are poorly structured and act to severely impede drainage as evidenced by gleying below and within the upper subsoils. Consequently these profiles are assigned to Wetness Class IV and are typified by Pit 1. The interaction between these heavy topsoils and drainage status with the climatic conditions which prevail at this site means that this land can be classified as no better than Subgrade 3b. This land may be subject to significantly reduced flexibility of cropping, stocking and cultivations.

ADAS Ref 2007/197/94  
MAFF Ref EL 20/328

Resource Planning Team  
Guildford Statutory Group  
ADAS Reading

## **SOURCES OF REFERENCE**

Geological Survey (1976) Sheet No 288 Maidstone 1 50 000 Series (solid and drift edition)

MAFF (1988) Agricultural Land Classification of England and Wales Revised guidelines and criteria for grading the quality of agricultural land

Meteorological Office (1989) Climatological Data for Agricultural Land Classification

Soil Survey of England and Wales (1983) Sheet 6 Soils of South East England 1 250 000 and accompanying legend

## APPENDIX I

### DESCRIPTION OF THE GRADES AND SUBGRADES

#### **Grade 1 Excellent Quality Agricultural Land**

Land with no or very minor limitations to agricultural use. A very wide range of agricultural and horticultural crops can be grown and commonly includes top fruit, soft fruit, salad crops and winter harvested vegetables. Yields are high and less variable than on land of lower quality.

#### **Grade 2 Very Good Quality Agricultural Land**

Land with minor limitations which affect crop yield, cultivations or harvesting. A wide range of agricultural or horticultural crops can usually be grown but on some land of this grade there may be reduced flexibility due to difficulties with the production of the more demanding crops such as winter harvested vegetables and arable root crops. The level of yield is generally high but may be lower or more variable than Grade 1 land.

#### **Grade 3 Good to Moderate Quality Land**

Land with moderate limitations which affect the choice of crops, the timing and type of cultivation, harvesting or the level of yield. When more demanding crops are grown, yields are generally lower or more variable than on land in Grades 1 and 2.

#### **Subgrade 3a Good Quality Agricultural Land**

Land capable of consistently producing moderate to high yields of a narrow range of arable crops, especially cereals, or moderate yields of a wide range of crops including cereals, grass, oilseed rape, potatoes, sugar beet and the less demanding horticultural crops.

#### **Subgrade 3b Moderate Quality Agricultural Land**

Land capable of producing moderate yields of a narrow range of crops, principally cereals and grass, or lower yields of a wider range of crops or high yields of grass which can be grazed or harvested over most of the year.

#### **Grade 4 Poor Quality Agricultural Land**

Land with severe limitations which significantly restrict the range of crops and/or the level of yields. It is mainly suited to grass with occasional arable crops (eg cereals and forage crops) the yields of which are variable. In most climates, yields of grass may be moderate to high but there may be difficulties in utilisation. The grade also includes very droughty arable land.

#### **Grade 5 Very Poor Quality Agricultural Land**

Land with severe limitations which restrict use to permanent pasture or rough grazing, except for occasional pioneer forage crops.

## **Urban**

Built-up or 'hard' uses with relatively little potential for a return to agriculture including housing industry commerce education, transport religious buildings cemeteries Also hard-surfaced sports facilities permanent caravan sites and vacant land all types of derelict land including mineral workings which are only likely to be reclaimed using derelict land grants

## **Non-agricultural**

'Soft' uses where most of the land could be returned relatively easily to agriculture including private parkland public open spaces sports fields allotments and soft surfaced areas on airports Also active mineral workings and refuse tips where restoration conditions to soft' after-uses may apply

## **Woodland**

Includes commercial and non-commercial woodland A distinction may be made as necessary between farm and non farm woodland

## **Agricultural Buildings**

Includes the normal range of agricultural buildings as well as other relatively permanent structures such as glasshouses Temporary structures (eg polythene tunnels erected for lambing) may be ignored

## **Open Water**

Includes lakes ponds and rivers as map scale permits

## **Land Not Surveyed**

Agricultural land which has not been surveyed

Where the land use includes more than one of the above eg buildings in large grounds and where map scale permits, the cover types may be shown separately Otherwise the most extensive cover type will be shown

## APPENDIX II

### FIELD ASSESSMENT OF SOIL WETNESS CLASS

#### SOIL WETNESS CLASSIFICATION

Soil wetness is classified according to the depth and duration of waterlogging in the soil profile. Six soil wetness classes are identified and are defined in the table below.

#### Definition of Soil Wetness Classes

Wetness Class	Duration of Waterlogging <sup>1</sup>
I	The soil profile is not wet within 70 cm depth for more than 30 days in most years <sup>2</sup>
II	The soil profile is wet within 70 cm depth for 31-90 days in most years or, if there is no slowly permeable layer within 80 cm depth it is wet within 70 cm for more than 90 days but only wet within 40 cm depth for 30 days in most years
III	The soil profile is wet within 70 cm depth for 91-180 days in most years or if there is no slowly permeable layer present within 80 cm depth it is wet within 70 cm for more than 180 days but only wet within 40 cm depth for between 31-90 days in most years
IV	The soil profile is wet within 70 cm depth for more than 180 days but not wet within 40 cm depth for more than 210 days in most years or if there is no slowly permeable layer present within 80 cm depth it is wet within 40 cm depth for 91-210 days in most years
V	The soil profile is wet within 40 cm depth for 211-335 days in most years
VI	The soil profile is wet within 40 cm depth for more than 335 days in most years

Soils can be allocated to a wetness class on the basis of quantitative data recorded over a period of many years or by the interpretation of soil profile characteristics, site and climatic factors. Adequate quantitative data will rarely be available for ALC surveys and therefore the interpretative method of field assessment is used to identify soil wetness class in the field. The method adopted here is common to ADAS and the SSLRC.

<sup>1</sup>The number of days specified is not necessarily a continuous period

<sup>2</sup>In most years is defined as more than 10 out of 20 years



**APPENDIX III**  
**SOIL PIT AND SOIL BORING DESCRIPTIONS**

**Contents**

**Soil Abbreviations - Explanatory Note**

**Soil Pit Descriptions**

**Database Printout - Boring Level Information**

**Database Printout - Horizon Level Information**

## SOIL PROFILE DESCRIPTIONS EXPLANATORY NOTE

Soil pit and auger boring information collected during ALC fieldwork is held on a computer database. This uses notations and abbreviations as set out below.

### Boring Header Information

- 1 **GRID REF** national 100 km grid square and 8 figure grid reference
- 2 **USE** Land use at the time of survey. The following abbreviations are used:

<b>ARA</b> Arable	<b>WHT</b> Wheat	<b>BAR</b> Barley
<b>CER</b> Cereals	<b>OAT</b> Oats	<b>MZE</b> Maize
<b>OSR</b> Oilseed rape	<b>BEN</b> Field Beans	<b>BRA</b> Brassicae
<b>POT</b> Potatoes	<b>SBT</b> Sugar Beet	<b>FCD</b> Fodder Crops
<b>LIN</b> Linseed	<b>FRT</b> Soft and Top Fruit	<b>FLW</b> Fallow
<b>PGR</b> Permanent Pasture	<b>LEY</b> Ley Grass	<b>RGR</b> Rough Grazing
<b>SCR</b> Scrub	<b>CFW</b> Coniferous Woodland	<b>DCW</b> Deciduous Wood
<b>HTH</b> Heathland	<b>BOG</b> Bog or Marsh	<b>FLW</b> Fallow
<b>PLO</b> Ploughed	<b>SAS</b> Set aside	<b>OTH</b> Other
<b>HRT</b> Horticultural Crops		
- 3 **GRDNT** Gradient as estimated or measured by a hand held optical clinometer
- 4 **GLEYSPL** Depth in centimetres (cm) to gleying and/or slowly permeable layers
- 5 **AP (WHEAT/POTS)** Crop adjusted available water capacity
- 6 **MB (WHEAT/POTS)** Moisture Balance (Crop adjusted AP crop adjusted MD)
- 7 **DRT** Best grade according to soil droughtiness
- 8 If any of the following factors are considered significant 'Y' will be entered in the relevant column:

<b>MREL</b> Microrelief limitation	<b>FLOOD</b> Flood risk	<b>EROSN</b> Soil erosion risk
<b>EXP</b> Exposure limitation	<b>FROST</b> Frost prone	<b>DIST</b> Disturbed land
<b>CHEM</b> Chemical limitation		
- 9 **LIMIT** The main limitation to land quality. The following abbreviations are used:

<b>OC</b> Overall Climate	<b>AE</b> Aspect	<b>EX</b> Exposure
<b>FR</b> Frost Risk	<b>GR</b> Gradient	<b>MR</b> Microrelief
<b>FL</b> Flood Risk	<b>TX</b> Topsoil Texture	<b>DP</b> Soil Depth
<b>CH</b> Chemical	<b>WE</b> Wetness	<b>WK</b> Workability
<b>DR</b> Drought	<b>ER</b> Erosion Risk	<b>WD</b> Soil Wetness/Droughtiness
<b>ST</b> Topsoil Stoniness		

## Soil Pits and Auger Borings

- 1 **TEXTURE** soil texture classes are denoted by the following abbreviations

<b>S</b>	Sand	<b>LS</b>	Loamy Sand	<b>SL</b>	Sandy Loam
<b>SZL</b>	Sandy Silt Loam	<b>CL</b>	Clay Loam	<b>ZCL</b>	Silty Clay Loam
<b>ZL</b>	Silt Loam	<b>SCL</b>	Sandy Clay Loam	<b>C</b>	Clay
<b>SC</b>	Sandy Clay	<b>ZC</b>	Silty Clay	<b>OL</b>	Organic Loam
<b>P</b>	Peat	<b>SP</b>	Sandy Peat	<b>LP</b>	Loamy Peat
<b>PL</b>	Peaty Loam	<b>PS</b>	Peaty Sand	<b>MZ</b>	Marine Light Silts

For the sand loamy sand sandy loam and sandy silt loam classes the predominant size of sand fraction will be indicated by the use of the following prefixes

<b>F</b>	Fine (more than 66% of the sand less than 0.2mm)
<b>M</b>	Medium (less than 66% fine sand and less than 33% coarse sand)
<b>C</b>	Coarse (more than 33% of the sand larger than 0.6mm)

The clay loam and silty clay loam classes will be sub-divided according to the clay content **M** Medium (<27% clay) **H** Heavy (27-35% clay)

- 2 **MOTTLE COL** Mottle colour using Munsell notation

- 3 **MOTTLE ABUN** Mottle abundance expressed as a percentage of the matrix or surface described

**F** few <2% **C** common 2-20% **M** many 20-40% **VM** very many 40% +

- 4 **MOTTLE CONT** Mottle contrast

**F** faint indistinct mottles evident only on close inspection  
**D** distinct - mottles are readily seen  
**P** prominent mottling is conspicuous and one of the outstanding features of the horizon

- 5 **PED COL** Ped face colour using Munsell notation

- 6 **GLEYS** If the soil horizon is gleyed a 'Y' will appear in this column. If slightly gleyed an 'S' will appear

- 7 **STONE LITH** Stone Lithology - One of the following is used

<b>HR</b>	all hard rocks and stones	<b>SLST</b>	soft oolitic or dolimitic limestone
<b>CH</b>	chalk	<b>FSST</b>	soft fine grained sandstone
<b>ZR</b>	soft argillaceous or silty rocks	<b>GH</b>	gravel with non-porous (hard) stones
<b>MSST</b>	soft medium grained sandstone	<b>GS</b>	gravel with porous (soft) stones
<b>SI</b>	soft weathered igneous/metamorphic rock		

Stone contents (>2cm, >6cm and total) are given in percentages (by volume)

8 **STRUCT** the degree of development size and shape of soil peds are described using the following notation

degree of development    **WK** weakly developed            **MD** moderately developed  
   **ST** strongly developed

ped size                            **F** fine                                    **M** medium  
   **C** coarse                                **VC** very coarse

ped shape                        **S** single grain                        **M** massive  
   **GR** granular                         **AB** angular blocky  
   **SAB** sub angular blocky        **PR** prismatic  
   **PL** platy

9 **CONSIST** Soil consistence is described using the following notation

**L** loose    **VF** very friable    **FR** friable    **FM** firm    **VM** very firm  
**EM** extremely firm            **EH** extremely hard

10 **SUBS STR** Subsoil structural condition recorded for the purpose of calculating profile droughtiness    **G** good    **M** moderate    **P** poor

11 **POR** Soil porosity If a soil horizon has less than 0.5% biopores >0.5 mm a 'Y' will appear in this column

12 **IMP** If the profile is impenetrable to rooting a 'Y' will appear in this column at the appropriate horizon

13 **SPL** Slowly permeable layer If the soil horizon is slowly permeable a 'Y' will appear in this column

14 **CALC** If the soil horizon is calcareous a 'Y' will appear in this column

15 Other notations

**APW** available water capacity (in mm) adjusted for wheat  
**APP** available water capacity (in mm) adjusted for potatoes  
**MBW** moisture balance wheat  
**MBP** moisture balance potatoes

SOIL PIT DESCRIPTION

Site Name MAIDSTONE LP SITE 46 (2) Pit Number 1P

Grid Reference TQ75374959 Average Annual Rainfall 681 mm  
 Accumulated Temperature 1447 degree days  
 Field Capacity Level 138 days  
 Land Use  
 Slope and Aspect 02 degrees S

HORIZON	TEXTURE	COLOUR	STONES >2	TOT STONE	LITH	MOTTLES	STRUCTURE	CONSIST	SUBSTRUCTURE	CALC
0- 30	HCL	10YR52 00	0	2	HR					
30- 36	HZCL	10YR63 00	0	2	HR	F	MDCSAB	FM	M	
36- 65	C	10YR63 00	0	2	HR	M	WKCSAB	FM	P	

Wetness Grade 3B Wetness Class IV  
 Gleying 036 cm  
 SPL 036 cm

Drought Grade APW mm MBW 0 mm  
 APP mm MBP 0 mm

FINAL ALC GRADE 3B  
 MAIN LIMITATION Wetness

SAMPLE NO	GRID REF	USE	ASPECT		--WETNESS--				-WHEAT-		-POTS-		M REL		EROSN	FROST	CHEM	ALC	COMMENTS	
			GRDNT	GLEY	SPL	CLASS	GRADE	AP	MB	AP	MB	DRT	FLOOD	EXP	DIST	LIMIT				
1	TQ75374959	ORC	S	02	028	036	4	3B		0	0							WE	3B	
1P	TQ75374959	ORC	S	02	036	036	4	3B		0	0							WE	3B	Pit dug to 65
2	TQ75344965	ORC	SW	02	027		2	3A		0	0							WE	3B	Imp 40 Re AB4
3	TQ75454958	ORC	S	01	028	028	4	3B		0	0							WE	3B	
4	TQ75354965	ORC	SW	02	030	045	3	3B		0	0							WE	3B	

SAMPLE	DEPTH	TEXTURE	COLOUR	----MOTTLES-----			PED COL	----STONES-----			STRUCT/ CONSIST	SUBS			SPL	CALC	
				COL	ABUN	CONT		GLE	>2	>6		LITH	TOT	STR			POR
1	0-28	hc1	10YR52 00						0	0	HR	2					
	28-36	hzc1	10YR53 00	10YR56	52	C			Y	0	0	HR	2		M		
	36-70	c	10YR63 00	10YR68	72	M	00MN00	00	Y	0	0	HR	5		P	Y	
1P	0-30	hc1	10YR52 00							0	0	HR	2				
	30-36	hzc1	10YR63 00	10YR56	00	F				0	0	HR	2	MDCSAB	FM	M	Y
	36-65	c	10YR63 00	10YR56	72	M	00MN00	00	Y	0	0	HR	2	WKCSAB	FM	P	Y
2	0-27	hzc1	10YR52 00							0	0	HR	2				
	27-40	hzc1	10YR63 00	10YR56	00	C	00MN00	00	Y	0	0	HR	5		M		
3	0-28	hzc1	10YR53 00	10YR56	00	F				0	0	HR	2				
	28-60	c	10YR53 00	10YR68	00	M			Y	0	0	HR	7		P	Y	
4	0-30	hzc1	10YR52 00							0	0	HR	2				
	30-45	hzc1	10YR63 00	10YR68	00	M	00MN00	00	Y	0	0	HR	5		M		
	45-60	c	10YR63 00	10YR68	52	M			Y	0	0	HR	5		P	Y	