

EC Directive 79/409 on the Conservation of Wild Birds Special Protection Area (SPA)

Name: Stour and Orwell Estuaries

Unitary Authority/County: Essex, Suffolk.

Site description: The Stour and Orwell estuaries straddle the eastern part of the Essex/Suffolk border in eastern England. The SPA is coincident with Cattawade Marshes Site of Special Scientific Interest (SSSI), Orwell Estuary SSSI and Stour Estuary SSSI. The estuaries include extensive mud-flats, low cliffs, saltmarsh and small areas of vegetated shingle on the lower reaches. The mud-flats hold *Enteromorpha*, *Zostera* and *Salicornia* spp. The site also includes areas of low-lying grazing marsh at Shotley Marshes on the south side of the Orwell and at Cattawade Marshes at the head of the Stour. Trimley Marshes on the north side of the Orwell includes several shallow freshwater pools, as well as areas of grazing marsh, and is managed as a nature reserve by the Suffolk Wildlife Trust. In summer, the site supports important numbers of breeding avocet *Recurvirostra avosetta*, while in winter it holds major concentrations of waterbirds, especially geese, ducks and waders. The geese also feed, and some waders roost, in surrounding areas of agricultural land outside the SPA. The site has close ecological links with the Hamford Water and Mid-Essex Coast SPAs, lying to the south on the same coast.

Size of SPA: The SPA covers an area of 3,676.92 ha.

Qualifying species:

The site qualifies under **article 4.1** of the Directive (79/409/EEC) as it is used regularly by 1% or more of the Great Britain populations of the following species listed in Annex I in any season:

Annex 1 species	Count and season	Period	% of GB population
Avocet <i>Recurvirostra avosetta</i>	21 pairs - breeding	5 year peak mean 1996 – 2000	3.6%

The site qualifies under **article 4.2** of the Directive (79/409/EEC) as it is used regularly by 1% or more of the biogeographical populations of the following regularly occurring migratory species (other than those listed in Annex I) in any season:

Migratory species	Count and season	Period	% of subspecies/population
Redshank <i>Tringa totanus</i>	2,588 individuals – autumn passage	5 year peak mean 1995/96 – 1999/2000	2.0% <i>britannica</i>
Dark-bellied brent goose <i>Branta bernicla bernicla</i>	2,627 individuals - wintering	5 year peak mean 1995/96 – 1999/2000	1.2% <i>bernicla</i> , Western Siberia (breeding)
Pintail <i>Anas acuta</i>	741 individuals - wintering	5 year peak mean 1995/96 – 1999/2000	1.2% Northwestern Europe (non-breeding)
Grey plover <i>Pluvialis squatarola</i>	3,261 individuals - wintering	5 year peak mean 1995/96 – 1999/2000	1.3% Eastern Atlantic (non- breeding)
Knot <i>Calidris canutus</i> <i>islandica</i>	5,970 individuals - wintering	5 year peak mean 1995/96 – 1999/2000	1.3% <i>islandica</i>
Dunlin <i>Calidris alpina alpina</i>	19,114 individuals - wintering	5 year peak mean 1995/96 – 1999/2000	1.4% <i>alpina</i> , Western Europe (non-breeding)
Black-tailed godwit <i>Limosa limosa islandica</i>	2,559 individuals - wintering	5 year peak mean 1995/96 – 1999/2000	7.3% <i>islandica</i>
Redshank <i>Tringa totanus</i>	3,687 individuals - wintering	5 year peak mean 1995/96 – 1999/2000	2.8% <i>britannica</i>

Bird counts from: Wetland Bird Survey (WeBS) database.

Assemblage qualification:

The site qualifies under **article 4.2** of the Directive (79/409/EEC) as it is used regularly by over 20,000 waterbirds (waterbirds as defined by the Ramsar Convention) in any season:

In the non-breeding season, the area regularly supports 63,017 individual waterbirds (5 year peak mean 1993/94 - 1997/98), including great crested grebe *Podiceps cristatus*, cormorant *Phalacrocorax carbo*, dark-bellied brent goose *Branta bernicla bernicla*, shelduck *Tadorna tadorna*, wigeon *Anas penelope*, gadwall *Anas strepera*, pintail *Anas acuta*, goldeneye *Bucephala clangula*, ringed plover *Charadrius hiaticula*, grey plover *Pluvialis squatarola*, lapwing *Vanellus vanellus*, knot *Calidris canutus islandica*, dunlin *Calidris alpina alpina*, black-tailed godwit *Limosa limosa islandica*, curlew *Numenius arquata*, redshank *Tringa totanus* and turnstone *Arenaria interpres*.

Non-qualifying species of interest: The SPA/Ramsar site as a whole, including the proposed extensions, is used by non-breeding marsh harrier *Circus aeruginosus*, hen harrier *Circus cyaneus*, merlin *Falco columbarius*, peregrine *Falco peregrinus*, short-eared owl *Asio flammeus* and kingfisher *Alcedo atthis* (all species listed in Annex I of the EC Birds Directive) in numbers of less than European importance (less than 1% GB population). It also supports breeding common tern *Sterna hirundo*, little tern *Sterna albifrons* and kingfisher (all listed in Annex I) in numbers of less than European importance.

Status of SPA:

- 1) Stour and Orwell Estuaries was classified as a Special Protection Area on 13 July 1994.
- 2) Extensions to the Stour and Orwell Estuaries SPA were classified on 19 May 2005.