

A1
Arun District Local Plan
Site 26 : Land south of
Findon Cemetery
Agricultural Land Classification
ALC Map and Report
April 1994

AGRICULTURAL LAND CLASSIFICATION REPORT

ARUN DISTRICT LOCAL PLAN

SITE 26 : LAND SOUTH OF FINDON CEMETERY

1. Summary

- 1.1 ADAS was commissioned by MAFF's Land Use Planning Unit to provide information on land quality for a number of sites in the Arun District of West Sussex. The work forms part of MAFF's statutory input to the preparation of the Arun District Local Plan.
- 1.2 Site 26 comprises 2 hectares of land to the south of Findon Cemetery, West Sussex. An Agricultural Land Classification, (ALC), survey was carried out during April 1994. The survey was undertaken at a detailed level of approximately one boring per hectare. A total of 3 borings and one soil inspection pit were described in accordance with MAFF's revised guidelines and criteria for grading the quality of agricultural land, (MAFF, 1988). These guidelines provide a framework for classifying land according to the extent to which its physical or chemical characteristics impose a long term limitation on its use for agriculture.
- 1.3 At the time of the survey the land use on the site was cereal cropping.
- 1.4 The distribution of grades and subgrades is shown on the attached ALC map and the areas and extent are given in the table below. The map has been drawn at a scale of 1:5,000. It is accurate at this scale, but any enlargement would be misleading.

Table 1 : Distribution of Grades and Subgrades

Grade	Area (ha)	% of Agricultural Land
2	0.3	15.0
3a	<u>1.7</u>	<u>85.0</u>
Total area of site	2.0	100% (1.0 ha)

- 1.5 Appendix I gives a general description of the grades, subgrades and land use categories identified in the survey. The main classes are described in terms of the type of limitation that can occur, the typical cropping range and the expected level and consistency of yield.
- 1.6 The majority of the site has been classified as good quality, Subgrade 3a land with a small area of very good quality, Grade 2, in the north. Soil droughtiness was shown to be the key limitation across the site. The Subgrade 3a profiles comprise medium silty clay loam topsoils, with large flints, over increasingly heavy and flinty subsoils which quickly became impenetrable. The Grade 2 profiles, on the other hand, comprise similar topsoils over chalky drift. The lower subsoils grow increasingly heavy and chalky before reaching the chalk bedrock at depth. The proportion of stones in both soil types reduce the amount of profile available water

for crops so adversely affecting the level and consistency of yield. The flinty soil represent a slightly greater drought risk than the chalky profiles, however, thus reducing land quality in this area.

2. Climate

- 2.1 The climatic criteria are considered first when classifying land as climate can be overriding in the sense that severe limitations will restrict land to low grades irrespective of favourable site or soil conditions.
- 2.2 The main parameters used in the assessment of an overall climatic limitation are average annual rainfall, as a measure of overall wetness, and accumulated temperature (degree days Jan-June), as a measure of the relative warmth of a locality.
- 2.3 A detailed assessment of the prevailing climate was made by interpolation from a 5km gridpoint dataset (Met. Office 1989). The details are given in the table below and these show that there is no overall climatic limitation affecting the site. However, climatic factors do interact with soil properties to influence soil wetness and droughtiness limitations.
- 2.4 No local climatic factors such as exposure or frost risk affect the site.

Table 2 : Climatic Interpolation

Grid Reference	TQ120086
Altitude (m)	57
Accumulated Temperature (days)	1477
Average Annual Rainfall (mm)	857
Field Capacity (days)	177
Moisture Deficit, Wheat (mm)	107
Moisture Deficit, Potatoes (mm)	100
Overall Climatic Grade	1

3. Relief

- 3.1 This site lies at the foot of a hill and slopes gently from 61m. AOD the north west to 52 m AOD in the south-east. Nowhere on the site do gradient or relief affect agricultural land quality.

4. Geology and Soil

- 4.1 British Geological Survey (1984), Sheet 318/333, Brighton and Worthing shows the site to be predominantly underlain by head deposits with a small area of Upper and Middle Chalk in the west.
- 4.2 The Soil Survey of England and Wales (1967) Sheets TQ00 and TQ10, show that the soils on this site comprise the Charity series in the south and the Coombe series in the north.. The Charity series is described as "well drained brown earths

developed in flinty silty head", (SSGB, 1967) while the Coombe series are "calcareous brown earths developed in silty, chalky head deposits". (SSGB, 1967).

- 4.3 Detailed field examination confirmed the existence of these soil types on the site but with just a thin band of Coombe in the north coincident with land mapped as Grade 2. The Upper and Middle Chalk, the parent material of the Coombe series, is obviously also found along the northern boundary contrary to the British Geological Survey's findings.

5. **Agricultural Land Classification**

- 5.1 Table 1 provided the details of the area measurements for each grade and the distribution of each grade is shown on the attached ALC map.

- 5.2 The location of the soil observation points are shown on the attached sample point map.

5.3 **Grade 2**

Along the northern boundary of the site a narrow band of very good quality agricultural land was identified. The profiles here comprise flinty (5% > 2 cm diameter, 1% flint > 6 cm diameter and 7% total flint v/v) calcareous medium silty clay loam topsoils over a similar textured upper subsoil containing 15% chalk v/v. In the heavy silty clay loam lower subsoil, between 40-65 cm from the surface, the chalk content markedly increases (25-60% total chalk v/v) before passing into the chalk bedrock at approximately 88 cm depth. It was considered that some water for crop growth would still be available in the top 30 cm of chalk but that the total amount of chalk and flint throughout the profile reduces the overall volume of water resulting in a slight drought risk. This marginally restricts the level and consistency of crop yield thus limiting the land to Grade 2 on soil droughtiness. The volume of topsoil flint, > 2 cm diameter, also imposes a limitation which is present in both soil units. Here farm machinery can become worn or damaged and the effective germination and growth of crops may be impeded. In addition, this land is limited to Grade 2 on the basis of topsoil workability. The presence of medium topsoil textures in a locality which is relatively moist, in a regional context, gives rise to a slight workability restriction.

5.4 **Subgrade 3a**

The remainder of the site has been graded as good quality agricultural land. The profiles comprise calcareous medium silty clay loam topsoils containing 8-9% flint > 2 cm diameter, 2% flint > 6 cm diameter and 10% total flint v/v. The upper subsoil varies in texture from medium or heavy silty clay loam to clay but becomes uniformly more flinty (10-15%) before reaching a layer, impenetrable to the auger, at 45-50 cm from the surface. Soil inspection pit 1 revealed this horizon to be clay

with 35% total flint v/v over a second clay horizon that continues to depth with 50% total flint. The high proportion of flint in this soil unit reduces the amount of profile available water for crops thus restricting level and consistency of yield and limiting the land to Subgrade 3a on soil droughtiness.

ADAS Ref: 4202/87/94
MAFF Ref: EL42/00460

Resource Planning Team
Guildford Statutory Group
ADAS Reading

SOURCES OF REFERENCE

British Geological Survey (1984), Sheet No. 318/333, Brighton and Worthing, 1:50,000

MAFF (1988), Agricultural Land Classification of England and Wales : Revised guidelines and criteria for grading the quality of agricultural land.

Meteorological Office (1989), Climatological Data for Agricultural Land Classification.

Soil Survey of England and Wales (1967), Sheet TQ00 and TQ10, Soil Maps of The West Sussex Coastal plain, 1:25,000.

Soil Survey of Great Britain (1967), Bulletin 3, Soils of the West Sussex Coastal Plain.

APPENDIX I

DESCRIPTION OF THE GRADES AND SUBGRADES

Grade 1 : Excellent Quality Agricultural Land

Land with no or very minor limitations to agricultural use. A very wide range of agricultural and horticultural crops can be grown and commonly includes top fruit, soft fruit, salad crops and winter harvested vegetables. Yields are high and less variable than on land of lower quality.

Grade 2 : Very Good Quality Agricultural Land

Land with minor limitations which affect crop yield, cultivations or harvesting. A wide range of agricultural or horticultural crops can usually be grown but on some land of this grade there may be reduced flexibility due to difficulties with the production of the more demanding crops such as winter harvested vegetables and arable root crops. The level of yield is generally high but may be lower or more variable than Grade 1 land.

Grade 3 : Good to Moderate Quality Land

Land with moderate limitations which affect the choice of crops, the timing and type of cultivation, harvesting or the level of yield. When more demanding crops are grown, yields are generally lower or more variable than on land in Grades 1 and 2.

Subgrade 3a : Good Quality Agricultural Land

Land capable of consistently producing moderate to high yields of a narrow range of arable crops, especially cereals, or moderate yields of a wide range of crops including cereals, grass, oilseed rape, potatoes, sugar beet and the less demanding horticultural crops.

Subgrade 3b : Moderate Quality Agricultural Land

Land capable of producing moderate yields of a narrow range of crops, principally cereals and grass, or lower yields of a wider range of crops or high yields of grass which can be grazed or harvested over most of the year.

Grade 4 : Poor Quality Agricultural Land

Land with severe limitations which significantly restrict the range of crops and/or the level of yields. It is mainly suited to grass with occasional arable crops (eg. cereals and forage crops) the yields of which are variable. In moist climates, yields of grass may be moderate to high but there may be difficulties in utilisation. The grade also includes very droughty arable land.

Grade 5 : Very Poor Quality Agricultural Land

Land with severe limitations which restrict use to permanent pasture or rough grazing, except for occasional pioneer forage crops.

Urban

Built-up or 'hard' uses with relatively little potential for a return to agriculture including: housing, industry, commerce, education, transport, religious buildings, cemeteries. Also, hard-surfaced sports facilities, permanent caravan sites and vacant land; all types of derelict land, including mineral workings which are only likely to be reclaimed using derelict land grants.

Non-agricultural

'Soft' uses where most of the land could be returned relatively easily to agriculture, including: private parkland, public open spaces, sports fields, allotments and soft-surfaced areas on airports. Also active mineral workings and refuse tips where restoration conditions to 'soft' after-uses may apply.

Woodland

Includes commercial and non-commercial woodland. A distinction may be made as necessary between farm and non-farm woodland.

Agricultural Buildings

Includes the normal range of agricultural buildings as well as other relatively permanent structures such as glasshouses. Temporary structures (eg. polythene tunnels erected for lambing) may be ignored.

Open Water

Includes lakes, ponds and rivers as map scale permits.

Land Not Surveyed

Agricultural land which has not been surveyed.

Where the land use includes more than one of the above, eg. buildings in large grounds, and where map scale permits, the cover types may be shown separately. Otherwise, the most extensive cover type will be shown.

APPENDIX II

FIELD ASSESSMENT OF SOIL WETNESS CLASS

SOIL WETNESS CLASSIFICATION

Soil wetness is classified according to the depth and duration of waterlogging in the soil profile. Six soil wetness classes are identified and are defined in the table below.

Definition of Soil Wetness Classes

Wetness Class	Duration of Waterlogging ¹
I	The soil profile is not wet within 70 cm depth for more than 30 days in most years. ²
II	The soil profile is wet within 70 cm depth for 31-90 days in most years or, if there is no slowly permeable layer within 80 cm depth, it is wet within 70 cm for more than 90 days, but only wet within 40 cm depth for 30 days in most years.
III	The soil profile is wet within 70 cm depth for 91-180 days in most years or, if there is no slowly permeable layer present within 80 cm depth, it is wet within 70 cm for more than 180 days, but only wet within 40 cm depth for between 31-90 days in most years.
IV	The soil profile is wet within 70 cm depth for more than 180 days but not wet within 40 cm depth for more than 210 days in most years or, if there is no slowly permeable layer present within 80 cm depth, it is wet within 40 cm depth for 91-210 days in most years.
V	The soil profile is wet within 40 cm depth for 211-335 days in most years.
VI	The soil profile is wet within 40 cm depth for more than 335 days in most years.

Soils can be allocated to a wetness class on the basis of quantitative data recorded over a period of many years or by the interpretation of soil profile characteristics, site and climatic factors. Adequate quantitative data will rarely be available for ALC surveys and therefore the interpretative method of field assessment is used to identify soil wetness class in the field. The method adopted here is common to ADAS and the SSLRC.

¹The number of days specified is not necessarily a continuous period.

²'In most years' is defined as more than 10 out of 20 years.

APPENDIX III

SOIL PIT AND SOIL BORING DESCRIPTIONS

Contents :

Soil Abbreviations - Explanatory Note

Soil Pit Descriptions

Database Printout - Boring Level Information

Database Printout - Horizon Level Information

SOIL PROFILE DESCRIPTIONS : EXPLANATORY NOTE

Soil pit and auger boring information collected during ALC fieldwork is held on a computer database. This uses notations and abbreviations as set out below.

Boring Header Information

1. **GRID REF** : national 100 km grid square and 8 figure grid reference.
2. **USE** : Land use at the time of survey. The following abbreviations are used.

ARA : Arable	WHT : Wheat	BAR : Barley
CER : Cereals	OAT : Oats	MZE : Maize
OSR : Oilseed rape	BEN : Field Beans	BRA : Brassicae
POT : Potatoes	SBT : Sugar Beet	FCD : Fodder Crops
LIN : Linseed	FRT : Soft and Top Fruit	FLW : Fallow
PGR : Permanent Pasture	LEY : Ley Grass	RGR : Rough Grazing
SCR : Scrub	CFW : Coniferous Woodland	DCW : Deciduous Wood
HTH : Heathland	BOG : Bog or Marsh	FLW : Fallow
PLO : Ploughed	SAS : Set aside	OTH : Other
HRT : Horticultural Crops		

3. **GRDNT** : Gradient as estimated or measured by a hand-held optical clinometer.
4. **GLEYSPL** : Depth in centimetres (cm) to gleying and/or slowly permeable layers.
5. **AP (WHEAT/POTS)** : Crop-adjusted available water capacity.
6. **MB (WHEAT/POTS)** : Moisture Balance. (Crop adjusted AP - crop adjusted MD)
7. **DRT** : Best grade according to soil droughtiness.
8. If any of the following factors are considered significant, 'Y' will be entered in the relevant column.

MREL : Microrelief limitation	FLOOD : Flood risk	EROSN : Soil erosion risk
EXP : Exposure limitation	FROST : Frost prone	DIST : Disturbed land
CHEM : Chemical limitation		

9. **LIMIT** : The main limitation to land quality. The following abbreviations are used.

OC : Overall Climate	AE : Aspect	EX : Exposure
FR : Frost Risk	GR : Gradient	MR : Microrelief
FL : Flood Risk	TX : Topsoil Texture	DP : Soil Depth
CH : Chemical	WE : Wetness	WK : Workability
DR : Drought	ER : Erosion Risk	WD : Soil Wetness/Droughtiness
ST : Topsoil Stoniness		

Soil Pits and Auger Borings

1. **TEXTURE** : soil texture classes are denoted by the following abbreviations.

S :	Sand	LS :	Loamy Sand	SL :	Sandy Loam
SZL :	Sandy Silt Loam	CL :	Clay Loam	ZCL :	Silty Clay Loam
ZL :	Silt Loam	SCL :	Sandy Clay Loam	C :	Clay
SC :	Sandy Clay	ZC :	Silty Clay	OL :	Organic Loam
P :	Peat	SP :	Sandy Peat	LP :	Loamy Peat
PL :	Peaty Loam	PS :	Peaty Sand	MZ :	Marine Light Silts

For the sand, loamy sand, sandy loam and sandy silt loam classes, the predominant size of sand fraction will be indicated by the use of the following prefixes:

F :	Fine (more than 66% of the sand less than 0.2mm)
M :	Medium (less than 66% fine sand and less than 33% coarse sand)
C :	Coarse (more than 33% of the sand larger than 0.6mm)

The clay loam and silty clay loam classes will be sub-divided according to the clay content: **M** : Medium (<27% clay) **H** : Heavy (27-35% clay)

2. **MOTTLE COL** : Mottle colour using Munsell notation.
3. **MOTTLE ABUN** : Mottle abundance, expressed as a percentage of the matrix or surface described.

F : few <2% **C** : common 2-20% **M** : many 20-40% **VM** : very many 40% +

4. **MOTTLE CONT** : Mottle contrast

F : faint - indistinct mottles, evident only on close inspection
D : distinct - mottles are readily seen
P : prominent - mottling is conspicuous and one of the outstanding features of the horizon

5. **PED. COL** : Ped face colour using Munsell notation.
6. **GLEYS** : If the soil horizon is gleyed a 'Y' will appear in this column. If slightly gleyed, an 'S' will appear.
7. **STONE LITH** : Stone Lithology - One of the following is used.

HR :	all hard rocks and stones	SLST :	soft oolitic or dolimitic limestone
CH :	chalk	FSST :	soft, fine grained sandstone
ZR :	soft, argillaceous, or silty rocks	GH :	gravel with non-porous (hard) stones
MSST :	soft, medium grained sandstone	GS :	gravel with porous (soft) stones
SI :	soft weathered igneous/metamorphic rock		

Stone contents (>2cm, >6cm and total) are given in percentages (by volume).

8. **STRUCT** : the degree of development, size and shape of soil peds are described using the following notation:

degree of development **WK** : weakly developed **MD** : moderately developed
 ST : strongly developed

ped size **F** : fine **M** : medium
 C : coarse **VC** : very coarse

ped shape **S** : single grain **M** : massive
 GR : granular **AB** : angular blocky
 SAB : sub-angular blocky **PR** : prismatic
 PL : platy

9. **CONSIST** : Soil consistence is described using the following notation:

L : loose **VF** : very friable **FR** : friable **FM** : firm **VM** : very firm
EM : extremely firm **EH** : extremely hard

10. **SUBS STR** : Subsoil structural condition recorded for the purpose of calculating profile droughtiness : **G** : good **M** : moderate **P** : poor

11. **POR** : Soil porosity. If a soil horizon has less than 0.5% biopores >0.5 mm, a 'Y' will appear in this column.

12. **IMP** : If the profile is impenetrable to rooting a 'Y' will appear in this column at the appropriate horizon.

13. **SPL** : Slowly permeable layer. If the soil horizon is slowly permeable a 'Y' will appear in this column.

14. **CALC** : If the soil horizon is calcareous, a 'Y' will appear in this column.

15. Other notations

APW : available water capacity (in mm) adjusted for wheat
APP : available water capacity (in mm) adjusted for potatoes
MBW : moisture balance, wheat
MBP : moisture balance, potatoes

SAMPLE NO.	GRID REF	USE	ASPECT	GRDNT	GLEYSPL	--WETNESS--		-WHEAT-		-POTS-		M.REL		EROSN	FROST	CHEM	ALC	COMMENTS	
						CLASS	GRADE	AP	MB	AP	MB	DRT	FLOOD						EXP
1	TQ12000870	CER	E	03		1	2	136	28	113	12	2					WD	2	CHALKY
1P	TQ12050852	CER	SE	01		1	2	91	-17	98	-3	3A					DR	3A	IMP 80
2	TQ12000860	CER	SE	03		1	2	72	-36	72	-29	3B					DR	3A	IMP 45 SEE 1P
3	TQ12050852	CER	SE	01		1	2	80	-28	80	-21	3B					DR	3A	IMP 50 SEE 1P

SAMPLE	DEPTH	TEXTURE	COLOUR	----MOTTLES-----			PED		----STONES----			STRUCT/ CONSIST	SUBS					
				COL	ABUN	CONT	COL.	GLE	>2	>6	LITH		TOT	STR	POR	IMP	SPL	CALC
1	0-30	mzc1	10YR42 00						5	1	HR	7						Y
	30-38	mzc1	10YR44 54						0	0	CH	15		M				Y
	38-65	hzc1	10YR54 00						0	0	CH	25		M				Y
	65-88	hzc1	10YR64 00						0	0	CH	60		P				Y
	88-120	ch	10YR81 00						0	0		0		P				Y
1P	0-24	mzc1	10YR42 00						9	2	HR	10						Y
	24-40	mzc1	10YR44 54						0	0	HR	10	MDCSAB	FR	M			Y
	40-70	c	10YR54 56						0	0	HR	35	WKCSAB	FM	M			Y
	70-80	c	10YR56 00						0	0	HR	50			P			Y
2	0-29	mzc1	10YR42 00						8	2	HR	10						Y
	29-45	c	10YR44 54						0	0	HR	15		M				Y
3	0-28	mzc1	10YR42 00						9	2	HR	10						Y
	28-50	hzc1	10YR54 00						0	0	HR	15		M				Y

SOIL PIT DESCRIPTION

Site Name : ARUN LP, SITE 26

Pit Number : 1P

Grid Reference: TQ12050852 Average Annual Rainfall : 857 mm
 Accumulated Temperature : 1482 degree days
 Field Capacity Level : 177 days
 Land Use : Cereals
 Slope and Aspect : 01 degrees SE

HORIZON	TEXTURE	COLOUR	STONES >2	TOT.STONE	MOTTLES	STRUCTURE
0- 24	MZCL	10YR4/2 00	9	10		
24- 40	MZCL	10YR4/4 54	0	10		MDCSAB
40- 70	C	10YR5/4 56	0	35		WKCSAB
70- 80	C	10YR5/6 00	0	50		

Wetness Grade : 2 Wetness Class : I
 Gleying : cm
 SPL : cm

Drought Grade : 3A APW : 91 mm MBW : -17 mm
 APP : 98 mm MBP : -3 mm

FINAL ALC GRADE : 3A
 MAIN LIMITATION : Droughtiness