

# COTSWOLD DISTRICT LOCAL PLAN: BLEDDINGTON

## AGRICULTURAL LAND CLASSIFICATION

### Report of survey

#### 1. INTRODUCTION

Four and a half hectares of land at Bledington were graded using the Agricultural Land Classification (ALC) System in September 1992. The survey was carried out for MAFF as part of its statutory input to the draft consultation on the Cotswold District Local Plan.

The fieldwork was carried out by ADAS's Resource Planning Team (Taunton Statutory Unit) at a scale of 1:10,000 (approximately one sample point every hectare). The information is correct at the scale shown but any enlargement would be misleading. This survey supercedes the 1" to the mile ALC map of this area being at a more detailed level and carried out under the Revised Guidelines and Criteria for grading the quality of agricultural land (MAFF 1988). A total of 3 borings were examined.

The ALC provides a framework for classifying land according to the extent to which its physical or chemical characteristics impose long-term limitations on agricultural use. The grading takes account of the top 120cm of the soil profile. A description of the grades used in the ALC System can be found in the appendix.

The distribution of ALC grades identified in the survey area is detailed below and illustrated on the accompanying map.

**Table 1 Distribution of ALC grades: Bledington**

Grade	Area (ha)	% of Survey Area	% of Agricultural Land
1	3.3	71.7	75.0
3B	1.1	23.9	25.0
Urban	<u>0.2</u>	<u>4.4</u>	100% (4.4ha)
TOTAL	4.6	100%	

## 2. CLIMATE

The grade of the land is determined by the most limiting factor present. The overall climate is considered first because it can have an overriding influence on restricting land to lower grades despite other favourable conditions.

To assess any overall climatic limitation, estimates of important climatic variables were obtained for the site by interpolation from the 5km grid Met Office/Maff Database (Met Office/ MAFF/ SSLRC 1989). The parameters used for assessing climate are accumulated temperature, (a measure of the relative warmth of a locality) and average annual rainfall, (a measure of overall wetness). The results shown in Table 2 reveal that there is no overall climatic limitation.

No local climatic factors such as exposure were noted in the survey area. Climatic data on Field Capacity Days (FCD) and Moisture Deficits for wheat (MDW) and potatoes (MDP) are also shown. This data is used in assessing the soil wetness and droughtiness limitations referred to in Section 5.

**Table 2 Climatic Interpolations: Bledington**

Grid Reference	SP 227 244	SP 228 243
Height (m)	115	110
Accumulated Temperature (° days)	1381	1387
Average Annual Rainfall (mm)	732	732
Overall Climatic Grade	1	1
Field Capacity (Days)	166	166
Moisture Deficit, Wheat (mm)	93	94
Potatoes (mm)	81	82

## 3. RELIEF

The survey area gently slopes to the south with a 5m drop in height from 115 AOD to 110 AOD.

## 4. GEOLOGY AND SOILS

The area is underlain by river gravels of the the Third and Forth Terraces as shown on BGS sheet 218.

The soils across the site are well drained and become slightly stony with depth. The topsoils are medium clay loams. Over part of the site the soils become lighter in texture and in part become heavier. Part of the site in the south has been disturbed.

## **5. AGRICULTURAL LAND CLASSIFICATION**

The distribution of ALC grades identified in the survey area are detailed in Section 1 and are shown on the accompanying ALC map. The information is correct at the scale shown but any enlargement would be misleading.

### **Grade 1**

Most of the site is Grade 1. The soils are well drained with a topsoil of medium clay loam overlying a slightly stoney loamy subsoil. There are no limitations to downgrade the soils.

### **Subgrade 3b**

The area mapped as 3b has been used for extraction. It lies at a lower level than the surrounding land. The topsoil is a stoney heavy clay loam over an impenetrable stoney subsoil. The high stone content at shallow depth leads to a droughtiness limitation, downgrading the unit to 3b.

## **DESCRIPTION OF THE GRADES AND SUBGRADES**

### **Grade 1 - excellent quality agricultural land**

Land with no or very minor limitations to agricultural use. A very wide range of agricultural and horticultural crops can be grown and commonly include top fruit, soft fruit, salad crops and winter harvested vegetables. Yields are high and less variable than on land of lower quality.

### **Grade 2 - very good quality agricultural land**

Land with minor limitations which affect crop yield, cultivations or harvesting. A wide range of agricultural and horticultural crops can usually be grown but on some land in the grade there may be reduced flexibility due to difficulties with the production of the more demanding crops such as winter harvested vegetables and arable root crops. The level of yield is generally high but may be lower or more variable than Grade 1.

### **Grade 3 - good to moderate quality agricultural land**

Land with moderate limitations which affect the choice of crops, timing and type of cultivation, harvesting or the level of yield. Where more demanding crops are grown yields are generally lower or more variable than on land in Grades 1 and 2.

#### **Subgrade 3a - good quality agricultural land**

Land capable of consistently producing moderate to high yields of a narrow range of arable crops, especially cereals, or moderate yields of a wide range of crops including cereals, grass, oilseed rape, potatoes, sugar beet and the less demanding horticultural crops.

#### **Subgrade 3b - moderate quality agricultural land**

Land capable of producing moderate yields of a narrow range of crops, principally cereals and grass or lower yields of a wider range of crops or high yields of grass which can be grazed or harvested over most of the year.

### **Grade 4 - poor quality agricultural land**

Land with severe limitations which significantly restrict the range of crops and/or level of yields. It is mainly suited to grass with occasional arable crops (eg cereals and forage crops) the yields of which are variable. In most climates, yields of grass may be moderate to high but there may be difficulties in utilisation. The grade also includes very droughty arable land.

### **Grade 5 - very poor quality agricultural land**

Land with very severe limitations which restrict use to permanent pasture or rough grazing, except for occasional pioneer forage crops.

## Descriptions of other land categories used on ALC maps

### **Urban**

Built-up or 'hard' uses with relatively little potential for a return to agriculture including: housing, industry, commerce, education, transport, religious buildings, cemeteries. Also, hard-surfaced sports facilities, permanent caravan sites and vacant land; all types of derelict land, including mineral workings which are only likely to be reclaimed using derelict land grants.

### **Non-agricultural**

'Soft' uses where most of the land could be returned relatively easily to agriculture, including: private park land, public open spaces, sports fields, allotments and soft-surfaced areas on airports/airfields. Also active mineral workings and refuse tips where restoration conditions to 'soft' after-uses may apply.

### **Agricultural buildings**

Includes the normal range of agricultural buildings as well as other relatively permanent structures such as glasshouses. Temporary structures (eg polythene tunnels erected for lambing) may be ignored.

### **Open water**

Includes lakes, ponds and rivers as map scale permits.

### **Land not surveyed**

Agricultural land which has not been surveyed.

Where the land use includes more than one of the above land cover types, eg buildings in large grounds, and where map scale permits, the cover types may be shown separately. Otherwise, the most extensive cover type will usually be shown.

Source: MAFF (1988) Agricultural Land Classification of England and Wales  
(Revised guidelines and criteria for grading the quality of agricultural land) Alnwick.