

A1
Canterbury Local Plan
CAN 3: Thanington, Kent
Agricultural Land Classification,
ALC Map and Report.
August 1995

AGRICULTURAL LAND CLASSIFICATION REPORT

CANTERBURY DISTRICT LOCAL PLAN CAN 3: THANINGTON.

1. Summary

- 1.1 ADAS was commissioned by MAFF's Land Use Planning Unit to provide information on land quality for a number of sites in the Canterbury District of Kent. The work forms part of MAFF's statutory input to the preparation of the Canterbury District Local Plan.
- 1.2 The site comprises 7.8 hectares of land to the south-east of Thanington near Canterbury in Kent. An Agricultural Land Classification (ALC) survey was carried out during March 1995. The survey was undertaken at a detailed level of approximately one boring per hectare of agricultural land surveyed. A total of 9 borings and one soil inspection pit were described according to MAFF's revised guidelines and criteria for grading the quality of agricultural land (MAFF, 1988). These guidelines provide a framework for classifying land according to the extent to which its physical or chemical characteristics impose a long term limitation on its use for agriculture.
- 1.3 The work was carried out by members of the Resource Planning Team in the Guildford Statutory Group of ADAS.
- 1.4 At the time of the survey the land was under winter wheat. The Woodland shown on the site is mature and deciduous. The Non-agricultural areas are scrub and an unsurfaced, unfenced track. The Urban area is a concrete track.
- 1.5 The distribution of grades and subgrades is shown on the attached ALC map and the areas and extent are given in the table below. The map has been drawn at a scale of 1:10,000. It is accurate at this scale, but any enlargement would be misleading.

Table 1: Distribution of Grades and Subgrades

Grade	Area (ha)	% of Site
2	6.8	87.2
Urban	0.1	1.3
Non-Agricultural	0.6	7.7
Woodland	<u>0.3</u>	<u>3.8</u>
Total area of site	7.8ha	100.0

- 1.6 Appendix I gives a general description of the grades, subgrades and land use categories identified in the survey. The main classes are described in terms of the type of limitation that can occur, the typical cropping range and the expected level and consistency of yield.

- 1.7 The agricultural land at this site has been classified as Grade 2 (very good quality). Principal limitations to land quality include soil wetness and soil droughtiness. The soils in this area comprise slightly stony, light loamy and medium silty topsoils over medium silty and clayey subsoils. In the local climatic regime, soils of this nature slightly reduce profile available water. As such, there is a slight risk of drought stress affecting plant growth and yield. The heavier silty and clayey subsoils were found to be slowly permeable. This causes a slight drainage impedence and leads to a soil wetness limitation. Soil wetness affects plant growth and yield and reduces the opportunities for cultivations and/or grazing without causing structural damage to the soil.

2. Climate

- 2.1 The climatic criteria are considered first when classifying land as climate can be overriding in the sense that severe limitations will restrict land to low grades irrespective of favourable site or soil conditions.
- 2.2 The main parameters used in the assessment of an overall climatic limitation are average annual rainfall, as a measure of overall wetness, and accumulated temperature, as a measure of the relative warmth of a locality.
- 2.3 A detailed assessment of the prevailing climate was made by interpolation from a 5km gridpoint dataset (Met. Office, 1989). The details are given in the table below and these show that there is no overall climatic limitation affecting the site.
- 2.4 No local climatic factors such as exposure or frost risk are believed to affect the site. However, climatic and soil factors interact to influence soil wetness and droughtiness limitations.

Table 2: Climatic Interpolation

Grid Reference	TR137564
Altitude, (m, AOD)	30
Accumulated Temperature (day degrees C., Jan.-June)	1464
Average Annual Rainfall (mm)	681
Field Capacity Days	143
Moisture deficit, wheat (mm)	118
Moisture deficit, potatoes (mm)	114
Overall Climatic Grade	1

3. Relief

- 3.1 The site lies between approximately 25 and 40m AOD. Overall the site slopes gently from west to east. Nowhere on the site does slope affect agricultural land quality.

4. Geology and Soils

- 4.1 The published geological information (BGS, 1982), shows the majority of the site to be underlain by Head Brickearth as a drift deposit. Towards the extreme west of the site Cretaceous Upper Chalk is shown.
- 4.2 The published soils information (SSEW, 1980), shows the site to be underlain by argillic brown earth type soils from the Hamble, Bursledon and Woodnesborough series. The legend accompanying the map describes these as, 'silty soils in brickearth, associated with loamy soils in Thanet and Woolwich Beds. Free drainage generally, but with local slight impedance.' (SSEW, 1980). The soils encountered at this site were of this broad type.

5. Agricultural Land Classification

- 5.1 Paragraph 1.5 provides the details of the area measurements for each grade and the distribution of each grade is shown on the attached ALC map.
- 5.2 The location of the soil observation points are shown on the attached sample point map.

Grade 2

- 5.3 Land of very good quality has been mapped over all the agricultural area at this site. The principal limitation to land quality is soil droughtiness; occasionally, soil wetness is equally limiting. Soils at this site commonly comprise a very slightly to slightly stony (up to 8% total flints) medium sandy silt loam, occasionally medium silty clay loam topsoil. This passes to a stoneless to slightly stony (up to 10% total flints) medium sandy silt loam or a borderline medium silty clay loam / silt loam upper subsoil. As depth increases so the soil profiles examined become more variable. The lower subsoil horizons comprise stoneless to moderately stony (up to 30% total flints) medium sandy silt loam, medium silty clay loam, heavy silty clay loam and, occasionally, silty clay textures. These were often found to be gleyed and from the pit observation, IP, also slowly permeable. Because of this slight drainage impedance many of the profiles studied are classified as Wetness Class II. This is not significant over the majority of the site as the relatively dry climate in this area and the easily workable nature of the topsoil gives a Grade 1 classification on workability. However towards the west of the site the slowly permeable horizon occurs at a shallower depth such that Wetness Class III is appropriate. Therefore Grade 2 has been applied in this area as both soil wetness and soil droughtiness are equally limiting. Soil wetness slightly restricts land utilisation as well as affecting plant growth and yield.
- 5.4 The majority of the soil profiles examined at this site, in combination with the local climate, experience a reduced amount of plant available water such that Grade 2 has been applied over the site on the basis of a slight soil droughtiness limitation. Soil droughtiness affects plant growth and yield. Occasional observations on the site were of Grade 1 quality. However these better soils have not been mapped as they occur spasmodically. Pit 1, for example, is technically classified as a Grade 1

profile. Information from a nearby site (CAN 26, ADAS Reference 2002/056/95) suggest that there are local variations in subsoil textures and that it is appropriate to place all of this land in a single unit of Grade 2.

ADAS Ref: 2002/045/95
MAFF Ref: EL20/642

Resource Planning Team
Guildford Statutory Group
ADAS Reading

SOURCES OF REFERENCE

British Geological Survey (1982), Sheet 289, Canterbury, Solid & Drift Edition. 1:50,000

MAFF (1988), Agricultural Land Classification of England and Wales : Revised guidelines and criteria for grading the quality of agricultural land.

Meteorological Office (1989), Climatic datasets for Agricultural Land Classification.

Soil Survey of England and Wales (1980), Bulletin No. 9, Soils of Kent.

Soil Survey of England and Wales (1983), Sheet No.6, Soils of South-East England, 1:250,000, and accompanying legend.

Soil Survey of England and Wales (1984), Bulletin No.15, Soils and their use in South-East England.

APPENDIX I

DESCRIPTION OF THE GRADES AND SUBGRADES

Grade 1 : Excellent Quality Agricultural Land

Land with no or very minor limitations to agricultural use. A very wide range of agricultural and horticultural crops can be grown and commonly includes top fruit, soft fruit, salad crops and winter harvested vegetables. Yields are high and less variable than on land of lower quality.

Grade 2 : Very Good Quality Agricultural Land

Land with minor limitations which affect crop yield, cultivations or harvesting. A wide range of agricultural or horticultural crops can usually be grown but on some land of this grade there may be reduced flexibility due to difficulties with the production of the more demanding crops such as winter harvested vegetables and arable root crops. The level of yield is generally high but may be lower or more variable than Grade 1 land.

Grade 3 : Good to Moderate Quality Land

Land with moderate limitations which affect the choice of crops, the timing and type of cultivation, harvesting or the level of yield. When more demanding crops are grown, yields are generally lower or more variable than on land in Grades 1 and 2.

Subgrade 3a : Good Quality Agricultural Land

Land capable of consistently producing moderate to high yields of a narrow range of arable crops, especially cereals, or moderate yields of a wide range of crops including cereals, grass, oilseed rape, potatoes, sugar beet and the less demanding horticultural crops.

Subgrade 3b : Moderate Quality Agricultural Land

Land capable of producing moderate yields of a narrow range of crops, principally cereals and grass, or lower yields of a wider range of crops or high yields of grass which can be grazed or harvested over most of the year.

Grade 4 : Poor Quality Agricultural Land

Land with severe limitations which significantly restrict the range of crops and/or the level of yields. It is mainly suited to grass with occasional arable crops (eg. cereals and forage crops) the yields of which are variable. In moist climates, yields of grass may be moderate to high but there may be difficulties in utilisation. The grade also includes very droughty arable land.

Grade 5 : Very Poor Quality Agricultural Land

Land with severe limitations which restrict use to permanent pasture or rough grazing, except for occasional pioneer forage crops.

Urban

Built-up or 'hard' uses with relatively little potential for a return to agriculture including: housing, industry, commerce, education, transport, religious buildings, cemeteries. Also, hard-surfaced sports facilities, permanent caravan sites and vacant land; all types of derelict land, including mineral workings which are only likely to be reclaimed using derelict land grants.

Non-agricultural

'Soft' uses where most of the land could be returned relatively easily to agriculture, including: private parkland, public open spaces, sports fields, allotments and soft-surfaced areas on airports. Also active mineral workings and refuse tips where restoration conditions to 'soft' after-uses may apply.

Woodland

Includes commercial and non-commercial woodland. A distinction may be made as necessary between farm and non-farm woodland.

Agricultural Buildings

Includes the normal range of agricultural buildings as well as other relatively permanent structures such as glasshouses. Temporary structures (eg. polythene tunnels erected for lambing) may be ignored.

Open Water

Includes lakes, ponds and rivers as map scale permits.

Land Not Surveyed

Agricultural land which has not been surveyed.

Where the land use includes more than one of the above, eg. buildings in large grounds, and where map scale permits, the cover types may be shown separately. Otherwise, the most extensive cover type will be shown.

APPENDIX II

FIELD ASSESSMENT OF SOIL WETNESS CLASS

SOIL WETNESS CLASSIFICATION

Soil wetness is classified according to the depth and duration of waterlogging in the soil profile. Six soil wetness classes are identified and are defined in the table below.

Definition of Soil Wetness Classes

Wetness Class	Duration of Waterlogging ¹
I	The soil profile is not wet within 70 cm depth for more than 30 days in most years. ²
II	The soil profile is wet within 70 cm depth for 31-90 days in most years or, if there is no slowly permeable layer within 80 cm depth, it is wet within 70 cm for more than 90 days, but only wet within 40 cm depth for 30 days in most years.
III	The soil profile is wet within 70 cm depth for 91-180 days in most years or, if there is no slowly permeable layer present within 80 cm depth, it is wet within 70 cm for more than 180 days, but only wet within 40 cm depth for between 31-90 days in most years.
IV	The soil profile is wet within 70 cm depth for more than 180 days but not wet within 40 cm depth for more than 210 days in most years or, if there is no slowly permeable layer present within 80 cm depth, it is wet within 40 cm depth for 91-210 days in most years.
V	The soil profile is wet within 40 cm depth for 211-335 days in most years.
VI	The soil profile is wet within 40 cm depth for more than 335 days in most years.

Soils can be allocated to a wetness class on the basis of quantitative data recorded over a period of many years or by the interpretation of soil profile characteristics, site and climatic factors. Adequate quantitative data will rarely be available for ALC surveys and therefore the interpretative method of field assessment is used to identify soil wetness class in the field. The method adopted here is common to ADAS and the SSLRC.

¹The number of days specified is not necessarily a continuous period.

²'In most years' is defined as more than 10 out of 20 years.

APPENDIX III

SOIL PIT AND SOIL BORING DESCRIPTIONS

Contents :

Soil Abbreviations - Explanatory Note

Soil Pit Descriptions

Database Printout - Boring Level Information

Database Printout - Horizon Level Information

SOIL PROFILE DESCRIPTIONS : EXPLANATORY NOTE

Soil pit and auger boring information collected during ALC fieldwork is held on a computer database. This uses notations and abbreviations as set out below.

Boring Header Information

1. **GRID REF** : national 100 km grid square and 8 figure grid reference.
2. **USE** : Land use at the time of survey. The following abbreviations are used.

ARA : Arable	WHT : Wheat	BAR : Barley
CER : Cereals	OAT : Oats	MZE : Maize
OSR : Oilseed rape	BEN : Field Beans	BRA : Brassicae
POT : Potatoes	SBT : Sugar Beet	FCD : Fodder Crops
LIN : Linseed	FRT : Soft and Top Fruit	FLW : Fallow
PGR : Permanent Pasture	LEY : Ley Grass	RGR : Rough Grazing
SCR : Scrub	CFW : Coniferous Woodland	DCW : Deciduous Wood
HTH : Heathland	BOG : Bog or Marsh	FLW : Fallow
PLO : Ploughed	SAS : Set aside	OTH : Other
HRT : Horticultural Crops		
3. **GRDNT** : Gradient as estimated or measured by a hand-held optical clinometer.
4. **GLEYSPL** : Depth in centimetres (cm) to gleying and/or slowly permeable layers.
5. **AP (WHEAT/POTS)** : Crop-adjusted available water capacity.
6. **MB (WHEAT/POTS)** : Moisture Balance. (Crop adjusted AP - crop adjusted MD)
7. **DRT** : Best grade according to soil droughtiness.
8. If any of the following factors are considered significant, 'Y' will be entered in the relevant column.

MREL : Microrelief limitation **FLOOD** : Flood risk **EROSN** : Soil erosion risk
EXP : Exposure limitation **FROST** : Frost prone **DIST** : Disturbed land
CHEM : Chemical limitation

9. **LIMIT** : The main limitation to land quality. The following abbreviations are used.

OC : Overall Climate	AE : Aspect	EX : Exposure
FR : Frost Risk	GR : Gradient	MR : Microrelief
FL : Flood Risk	TX : Topsoil Texture	DP : Soil Depth
CH : Chemical	WE : Wetness	WK : Workability
DR : Drought	ER : Erosion Risk	WD : Soil Wetness/Droughtiness
ST : Topsoil Stoniness		

Soil Pits and Auger Borings

1. **TEXTURE** : soil texture classes are denoted by the following abbreviations.

S :	Sand	LS :	Loamy Sand	SL :	Sandy Loam
SZL :	Sandy Silt Loam	CL :	Clay Loam	ZCL :	Silty Clay Loam
ZL :	Silt Loam	SCL :	Sandy Clay Loam	C :	Clay
SC :	Sandy Clay	ZC :	Silty Clay	OL :	Organic Loam
P :	Peat	SP :	Sandy Peat	LP :	Loamy Peat
PL :	Peaty Loam	PS :	Peaty Sand	MZ :	Marine Light Silts

For the sand, loamy sand, sandy loam and sandy silt loam classes, the predominant size of sand fraction will be indicated by the use of the following prefixes:

F :	Fine (more than 66% of the sand less than 0.2mm)
M :	Medium (less than 66% fine sand and less than 33% coarse sand)
C :	Coarse (more than 33% of the sand larger than 0.6mm)

The clay loam and silty clay loam classes will be sub-divided according to the clay content: **M** : Medium (<27% clay) **H** : Heavy (27-35% clay)

2. **MOTTLE COL** : Mottle colour using Munsell notation.
3. **MOTTLE ABUN** : Mottle abundance, expressed as a percentage of the matrix or surface described.

F : few <2% **C** : common 2-20% **M** : many 20-40% **VM** : very many 40% +

4. **MOTTLE CONT** : Mottle contrast

F : faint - indistinct mottles, evident only on close inspection
D : distinct - mottles are readily seen
P : prominent - mottling is conspicuous and one of the outstanding features of the horizon

5. **PED. COL** : Ped face colour using Munsell notation.
6. **GLEYS** : If the soil horizon is gleyed a 'Y' will appear in this column. If slightly gleyed, an 'S' will appear.
7. **STONE LITH** : Stone Lithology - One of the following is used.

HR :	all hard rocks and stones	SLST :	soft oolitic or dolimitic limestone
CH :	chalk	FSST :	soft, fine grained sandstone
ZR :	soft, argillaceous, or silty rocks	GH :	gravel with non-porous (hard) stones
MSST :	soft, medium grained sandstone	GS :	gravel with porous (soft) stones
SI :	soft weathered igneous/metamorphic rock		

Stone contents (>2cm, >6cm and total) are given in percentages (by volume).

SOIL PIT DESCRIPTION

Site Name : CANTERBURY LP CAN 3 Pit Number : 1P

Grid Reference: TR13705650 Average Annual Rainfall : 681 mm
 Accumulated Temperature : 1464 degree days
 Field Capacity Level : 143 days
 Land Use : Cereals
 Slope and Aspect : degrees

HORIZON	TEXTURE	COLOUR	STONES >2	TOT. STONE	LITH	MOTTLES	STRUCTURE	CONSIST	SUBSTRUCTURE	CALC
0- 23	MSZL	10YR32 42	2	5	HR					
23- 73	ZL	10YR44 00	0	2	HR		MDCSAB	FR	M	
73-100	MZCL	10YR53 00	0	2	HR	C	MDCAB	FR	M	
100-120	HZCL	10YR53 00	0	10	HR	C		FR	M	

Wetness Grade : 1 Wetness Class : II
 Gleying : 73 cm
 SPL : 73 cm

Drought Grade : 1 APW : 176mm MBW : 58 mm
 APP : 143mm MBP : 29 mm

FINAL ALC GRADE : 1
 MAIN LIMITATION :

SAMPLE NO.	GRID REF	ASPECT USE	GRDNT	GLEYS	--WETNESS--		-WHEAT-		-POTS-		M.REL		EROSN	FROST		CHEM	ALC	COMMENTS	
					SPL	CLASS	GRADE	AP	MB	AP	MB	DRT	FLOOD	EXP	DIST	LIMIT			
1P	TR13705650	CER		73	73	2	1	176	58	143	29	1					1	PIT 100 @ AB 3	
2	TR13605650	CER				1	1	162	44	121	7	2				DR	2		
3	TR13705650	CER		70	70	2	1	179	61	145	31	1					1	SEE 1P	
4	TR13805650	SAS				1	1	159	41	123	9	2				DR	2		
5	TR13505640	CER		60	60	2	2	157	39	121	7	2				WD	2		
6	TR13605640	CER	N	02	80	80	2	1	158	40	122	8	2				DR	2	
7	TR13705640	CER			90	90	1	1	157	39	122	8	2				DR	2	
8	TR13805640	SAS				1	1	160	42	124	10	1					1		
9	TR13305630	CER	NW	02			1	1	109	-9	113	-1	3A				DR	2	IMP FLINTS 80
10	TR13425634	CER	NE	04	45	45	3	2	141	23	115	1	2				WD	2	

SAMPLE	DEPTH	TEXTURE	COLOUR	----MOTTLES-----			PED	----STONES-----			STRUCT/	SUBS						
				COL	ABUN	CONT	COL.	GLE	>2	>6	LITH	TOT	CONSIST	STR	POR	IMP	SPL	CALC
1P	0-23	msz1	10YR32 42					2	0	HR	5							
	23-73	z1	10YR44 00					0	0	HR	2	MDCSAB	FR	M				1% off mzc1
	73-100	mzc1	10YR53 00	10YR56 00	C	F		Y	0	0	HR	2	MDCAB	FR	M	Y		Y
	100-120	hzc1	10YR53 00	10YR56 00	C	F		Y	0	0	HR	10		FR	M	Y		Y
2	0-30	msz1	10YR32 00					0	0	HR	3							
	30-45	msz1	10YR44 54					0	0	HR	3			M				
	45-70	msz1	10YR54 00					0	0	HR	3			M				
	70-120	msz1	10YR54 00					0	0	HR	5			M				
3	0-20	msz1	10YR32 00					0	0	HR	3							
	20-70	z1	10YR44 00					0	0	HR	2			M				see 1p 1% off mzc1
	70-120	mzc1	10YR53 00	10YR56 00	C			Y	0	0	HR	1		M			Y	
4	0-27	msz1	10YR32 00					0	0	HR	2							
	27-45	msz1	10YR44 00					0	0	HR	2			M				
	45-120	mzc1	10YR54 00					0	0		0			M				
5	0-33	mzc1	10YR32 42					2	0	HR	8							
	33-60	mzc1	10YR54 00	10YR56 00	F			0	0		0			M				
	60-120	hzc1	10YR54 52	10YR56 00	C		00MN00	00	Y	0	0	0		M			Y	
6	0-25	msz1	10YR32 42					3	0	HR	5							
	25-80	mzc1	10YR54 00	10YR56 00	F			0	0		0			M				
	80-120	hzc1	10YR53 52	10YR56 00	C			Y	0	0	0			M			Y	
7	0-27	msz1	10YR32 00					0	0	HR	4							
	27-50	mzc1	10YR44 54					0	0	HR	1			M				
	50-90	mzc1	10YR54 00					0	0		0			M				
	90-120	hzc1	10YR53 00	10YR56 62	C		00MN00	00	Y	0	0	HR	4		M		Y	
8	0-30	msz1	10YR32 00					0	0	HR	1							
	30-50	mzc1	10YR54 00					0	0	HR	1			M				
	50-70	mzc1	10YR54 00					0	0		0			M				
	70-120	hzc1	10YR54 00					0	0		0			M				
9	0-30	msz1	10YR32 42					2	0	HR	8							
	30-65	mzc1	10YR54 00	10YR56 00	F			0	0	HR	10			M				
	65-80	msz1	10YR54 00	10YR56 00	F			0	0	HR	30			M				IMP FLINTS 80
10	0-28	msz1	10YR32 42					2	0	HR	5							
	28-45	mzc1	10YR44 54	00MN00 00	F			0	0	HR	5			M				
	45-120	zc	10YR53 00	10YR66 00	C			Y	0	0	HR	2		M		Y		