EC-Directive 79/409 on the Conservation of Wild Birds: Special Protection Area

SEVERN ESTUARY (GLOUCESTERSHIRE, AVON, SOMERSET, SOUTH GLAMORGAN, MID GLAMORGAN, GWENT)

The Severn Estuary is one of the largest estuaries in Britain and it has the second largest tidal range in the world. Its classic funnel shape and south-west orientation makes it susceptible to extreme weather conditions in the east Atlantic. There are large urban developments on the estuary including the cities of Bristol and Cardiff.

The Severn Estuary qualifies under Article 4.1 of the Birds Directive by regularly supporting an internationally important wintering population of Bewick's swan *Cygnus columbianus bewickii*, an Annex 1 species. During the period 1988/89 to 1992/93 a mean peak of 289 birds (1.7% of the northwest European population, 4.1% of the British wintering population) used the estuary.

The Severn Estuary qualifies under Article 4.2 as a wetland of international importance by regularly supporting in winter over 20,000 waterfowl. In the five year period 1988/89 to 1992/93 the average peak count was 68,026 waterfowl comprising 17,502 wildfowl and 50,524 waders.

The Severn Estuary also qualifies under Article 4.2 by regularly supporting in winter internationally important numbers of the following 5 species of migratory waterfowl (average peak means for the period 1988/89 to 1992/93): 3,002 European white-fronted goose *Anser albifrons albifrons* (1.0% NW European, 50.0% British), 2,892 shelduck *Tadorna tadorna* (1.2% NW European, 3.9% British), 330 gadwall *Anas strepera* (2.8% NW European, 5.5% British), 41,683 dunlin *Calidris alpina* (2.9% east Atlantic flyway (EAF), 9.6% British) and 2,013 redshank *Tringa totanus* (1.3% EAF, 2.6% British).

The Severn Estuary also supports nationally important wintering populations of a further 10 species: 3,977 wigeon *Anas penelope* (1.6% British), 1,998 teal *Anas crecca* (2.0% British), 523 pintail *Anas acuta* (2.1% British), 1,686 pochard *Aythya ferina* (3.8% British), 913 tufted duck *Aythya fuligula* (1.5% British), 227 ringed plover *Charadrius hiaticula* (1.0% British), 781 grey plover *Pluvialis squatarola* (3.7% British), 3,096 curlew *Numenius arquata* (3.4% British), 246 whimbrel N. phaeopus (4.9% British total) and 3 spotted redshank *Tringa erythropus* (1.5% British).

In addition, during passage periods, the estuary supports nationally important numbers of ringed plover (spring migration: 442 birds (1.4% British passage), autumn migration: 1,573 birds (5.2% British passage)), dunlin (spring: 3,510 birds (1.7% British passage), autumn: 5,500 birds (2.7% British passage)), whimbrel *Numenius phaeopus* (spring: 246 birds (4.9% British passage), autumn: 66 birds (1.3% British passage)) and redshank (autumn: 2,456 birds (2% British passage)).

The Severn Estuary also supports a nationally important breeding population of a migratory species. In 1993 2040 pairs of lesser black-backed gulls *Larus fuscus* bred on the islands of Steep Holm and Flat Holm within the estuary. This represents 2.5% of the British total.

SPA Citation CAR December 1993 This citation / map relates to a site entered in the Register of European sites for Great Britain Register reference number WOOLSONS.

Date of registration 2.5 AUS 1902

Signed 1 Register of the Environment